WHITE RIVER BASIN SURVEY: WEST FORK WHITE RIVER, 2004

Kevin J. Hoffman Assistant Research Biologist



Fisheries Section
Indiana Department of Natural Resources
Division of Fish and Wildlife
I.G.C.-South, Room W273
402 W. Washington Street
Indianapolis, IN 46204

EXECUTIVE SUMMARY

- Eighteen, approximately one mile reaches of river were sampled beginning at river mile 3.6 near the confluence with the Wabash River and ending at RM 194.1 near Martinsville, Indiana. Fish and habitat data were collected at each site. This report was part of a two-year project to assess fisheries data from the EFWR and the WFWR.
- There was no relation between IBI and QHEI scores. There was a positive correlation between IBI scores and river mile; IBI scores increased as river mile increased. The average IBI score was 40 (median = 40), which is considered "fair." Of 18 sampling reaches, 5 were "good" or better and the other 13 were scored as "fair" or lower. Fish habitat scores (QHEI) ranged from 44.0 to 66.0 (median = 56.8) for all reaches.
- All sampling efforts yielded 26,925 fish weighing approximately 3,970 lbs, representing 17 families, 70 species and two hybrids. Crews collected 55 species via electrofishing and an additional 17 species via seines.
- Spotfin shiner was the most abundant species (66%), followed by bullhead minnow (7%), sand shiner (6%), and bluntnose minnow (4%). Common carp (26%) was the most abundant species collected by weight, followed by smallmouth buffalo (15%), river carpsucker (10%), and blue sucker (9%). Sportfish comprised less than 2% of the total sample by number and less than 9% by weight.
- Channel catfish grew slower in the WFWR than in the EFWR. Flathead catfish and
 freshwater drum grew similarly in the EFWR and WFWR. Channel catfish up to 25.6
 and flathead catfish up to 33.6 in TL were collected. Spotted bass were slightly longer on
 average in the WFWR compared to the EFWR. Spotted bass up to 14.2 in TL were
 collected. Not enough data was collected from other sportfish species to do any
 comparisons.
- There were 19 public access sites for the entire WFWR. Only one site was recommended for acquisition by fisheries biologists. There were four access sites (Bloomfield, Freedom, Gosport, and Paragon) on the acquisition priority list as of February 2006.
- The only recommendation is to form a workgroup to develop a White River Basin management plan.

TABLE OF CONTENTS

	Page
INTRODUCTION	1
METHODS	1
RESULTS	2
Water chemistry and fish habitat	2
Fisheries survey data	2
DISCUSSION	7
RECOMMENDATIONS	8
LITERATURE CITED	9
TABLES	10
FIGURES	15
APPENDIX A	19
APPENDIX B	20
APPENDIX C	23
APPENDIX D	26
APPENDIX E	58
APPENDIX F	76

INTRODUCTION

The West Fork White River (WFWR) begins in Randolph County, in the east-central side of the state and flows 356 mi southwest and joins the East Fork White River in Pike and Daviess Counties. The WFWR drains approximately 5,372 mi². The upper reaches of the WFWR drain mostly agricultural fields until it flows through large cities, like Anderson and Indianapolis. Approximately 30 mi of the WFWR flows from the northeast corner of Indianapolis to the southwest corner of the city. There are multiple dams along the WFWR that impede fish from moving upriver, for example, the 16th St. Dam in southern Indianapolis and Broad Ripple Dam in northern Indianapolis. Commercial fishing is allowed on the WFWR from the mouth to the confluence with the Eel River.

The WFWR has experienced two large fish kills in the last 30 years. In 1976, a large fish kill affected almost 176 mi of river from Southport Rd. in Indianapolis to the mouth of the WFWR. An estimated 5 million fish were lost due to excess amounts of ammonia discharge into the river (Kingsley 1983). In December 1999, another major fish kill affected approximately 60 river mi from Anderson to Lake Indy. An estimated 4.3 million fish weighing 180 tons were lost after Guide Corporation used too much sodium dimethyldithiocarbamate (HMP 2000) to treat their wastewater that was eventually pumped directly into the city of Anderson's wastewater treatment facility. The HMP 2000 broke down into thiram, which is classified as a general use pesticide. The city's plant was quickly overcome and as a result, large quantities of contaminated wastewater and raw sewage were released into the WFWR.

Major portions of the West Fork White River have not been surveyed by the IDFW while some portions have been surveyed sporadically. The objective of this project was to collect fish community, fish habitat, and public access data to provide information needed to develop a riverwide management plan. Along with fish and public access data, recreational use data was collected in 2003 via a creel survey from April to October (Hoffman 2004).

METHODS

Eighteen, approximately one mi reaches of river were established beginning at river mi 3.6 and ending at RM 194.1 near Martinsville (Figure 1). Water chemistry data was collected at each reach according to Manual of Fisheries Survey Methods (Shipman 2001). Habitat data was

collected and scored at each reach according to the Qualitative Habitat Evaluation Index (QHEI) (Rankin 1989). Fish were sampled using boat-mounted DC electrofishing gear and sampling followed standard protocol for rivers and streams in Indiana (Shipman 2001). Each bank of the reach was sampled with approximately 0.5 h of effort. Additionally, a minimum of three seine hauls were conducted at each reach. All fish were measured to the nearest 0.1 in total length (TL). Fish weights were estimated using weight-length regressions. Scales were collected from a subsample of five fish per 0.5 in-group for sportfish. An IBI score was assigned to each reach based upon modifications by Simon (1998).

RESULTS

Water chemistry and fish habitat

Water chemistry data were within ranges to sustain fish survival (Table 1). Secchi disk measurements ranged from 9 to 21 in. Dissolved oxygen ranged from 8.0 to 17.0 ppm. Conductivity measurements ranged from 656 to 1,116 μ S. Water temperatures ranged from 67.8 to 80.6 °F. Average flow ranged from 837 to 3,400 cfs (Table 2). Gradient ranged from 0.4 to 2.84 ft/mi. Average width ranged from 187 to 506 ft. Fish habitat scores (QHEI) ranged from 44.0 to 66.0 (median = 56.8) for all reaches (Table 3).

Fisheries survey data

All sampling efforts yielded 26,925 fish weighing approximately 3,970 lbs, representing 17 families and 72 species and two hybrids. Crews collected 55 species via electrofishing and an additional 17 species via seines. Spotfin shiner was the most abundant species (66%), followed by bullhead minnow (7%), sand shiner (6%), and bluntnose minnow (4%). Common carp (26%) was the most abundant species collected by weight, followed by smallmouth buffalo (15%), river carpsucker (10%), and blue sucker (9%). Sportfish comprised less than 2% of the total sample by number and less than 9% by weight.

For combined electrofishing and seining data, IBI scores ranged from 32 to 46 of a maximum of 60 (Table 4). There was no significant correlation between QHEI and IBI scores for all reaches pooled (R = 0.2487; P = 0.320) (Figure 2). The average IBI score was 40 (median = 40), which is considered "fair." Of 18 sampling reaches, 5 were "good" or better and the other

13 were scored as "fair" or lower. There was a significant increase in IBI as river mi increased (R = 0.6748; P = 0.002) (Figure 2).

Carp and minnow family (Cyprinidae)

The Cyprinidae family comprised 21 species and accounted for 88% of the total number of fish collected and 29% of the total weight of fish collected. Spotfin shiner, bullhead minnow, sand shiner, and bluntnose minnow were among the most abundant species in this family. Spotfin shiner accounted for 66% of the total number of fish collected. Silver carp, silver chub, river shiner, bigeye chub, and channel shiner were among the least collected species from this family. Seining accounted for almost 98% of the individuals collected from this family.

Sucker family (Catostomidae)

The Catostomidae family comprised 12 species and represented 4% of the total sample by number, but 44% by weight. River carpsucker, smallmouth buffalo, blue sucker, and quillback were among the most abundant species of this family. Silver redhorse, black redhorse, and northern hogsucker were collected in fewer numbers. Only six northern hogsuckers were collected from only one sampling reach. Other species collected were golden redhorse, shorthead redhorse, highfin carpsucker, black buffalo, and bigmouth buffalo. Approximately 93% of the individuals from this family were collected via electrofishing.

Livebearer family (Poeciliidae)

The Poeciliidae family was represented by one species, the western mosquitofish. This species accounted for 3% of the total number of fish collected. Western mosquitofish were collected at 15 of 18 sampling reaches.

New World silversides (Atherinopsidae)

The only representative of this family was the brook silverside. This species (n = 318) accounted for just over 1% of the total sample by number and they were all collected by seining. Brook silverside ranged from 1.6 to 3.0 in TL and were collected at 11 of 18 sampling reaches.

Herring family (Clupeidae)

Gizzard shad and threadfin shad were the only representatives of the Clupeidae family and comprised 1% of the total sample by number and 2% by weight. Nearly all of the Clupeids were collected via electrofishing and they were mostly gizzard shad. Gizzard shad ranged from 2.3 to 16.8 in TL. Including only fish caught via electrofishing, gizzard shad accounted for 12% of the sample by number. Only one threadfin shad was collected at RM 57.5 and no gizzard shad were collected at this sampling reach.

Gar family (Lepisosteidae)

The Lepisosteidae family comprised two species and accounted for 1% of the total sample by number and nearly 9% by weight. Shortnose gar was more abundant than longnose gar. Shortnose gar ranged from 16.9 to 29.5 in TL and longnose gar ranged from 17.9 to 45 in TL. Gar were collected at all but one sampling reach.

Drum family (Sciaenidae)

Freshwater drum was the only representative of this family. Freshwater drum ranged from 3.7 to 24.1 in TL (Figure 3) and they accounted for less than 1% of the total sample by number and 6% of the total sample by weight. Freshwater drum were collected at all 18 sampling reaches. Freshwater drum reached 16.7 in by age 4. Some of the longest and potentially oldest freshwater drum were not aged because of regenerated scales.

Sunfish family (Centrarchidae)

The Centrarchidae family comprised 10 species and 1 hybrid, and represented less than 1% of the total sample by number and 1% by weight. Most of the Centrarchids were collected via electrofishing and they represented 7% of the electrofishing sample by number. Spotted bass was the most numerous sunfish species. Spotted bass were collected at 16 of 18 sampling reaches and they ranged from 2.2 to 14.2 in TL (Figure 3). Spotted bass reached 11.5 in TL by age 4 (Table 5). A von Bertalanffy growth function could not be fit to age data for spotted bass, presumably due to error in aging scales, especially in older fish.

Longear sunfish and bluegill were the next most common sunfish species. Longear sunfish up to 5.2 in TL and bluegill up to 6.0 in TL were collected. Bluegill were collected at 12 sampling reaches and longear were only collected at 9 sampling reaches.

White crappie, smallmouth bass, largemouth bass, black crappie, and redear sunfish were all caught in small numbers ($n \le 6$ each). White crappie were collected at five sampling reaches, smallmouth bass at two, largemouth bass and black crappie at one each. White crappie up to 12.8 in TL were collected. Green sunfish and orangespotted sunfish were also collected in few numbers.

Catfish family (Ictaluridae)

The Ictaluridae family comprised 4 species, accounting for less than 1% of the total number collected and 6% of the total weight collected. One individual blue catfish measuring 20.2 in TL and four mountain madtoms were collected. Channel catfish was the most abundant species in this family, accounting for almost 4% of the total weight collected. Channel catfish were collected at 16 of 18 sampling reaches and ranged from 1.8 to 25.6 in TL (Figure 3). The growth coefficient (K) from a von Bertalanffy growth curve was 0.138, which was lower than channel catfish in the EFWR (K = 0.165) (Hoffman 2006). The oldest individual was 12 years old (Table 5). Based on yield per recruit models using a natural mortality estimate of 25% and a total annual mortality estimate of 34% (based on EFWR channel catfish), channel catfish in the WFWR were not experiencing growth overfishing. If, however, exploitation were increased to 25% (about a 15% increase), growth overfishing would be occurring with a minimum size limit of 10 in TL.

Flathead catfish was the next most abundant species of the Ictaluridae family. Flathead catfish ranged from 4.0 to 33.1 in TL, accounting for 3% of the total sample. The majority of the flathead catfish were between 6 and 16 in TL. Not enough data was collected to fit a von Bertalannfy growth function to age data; however, flathead catfish reached 15.8 in TL by age 4 (Table 5).

Temperate bass family (Moronidae)

The Moronidae family consisted of three species and one hybrid. White bass was the most abundant species in this family and they ranged from 3.2 to 16.3 in TL. White bass (n =

35) were collected at 14 of the 18 sampling reaches. White bass up to age 6 were collected. Only six yellow bass, five hybrid striped bass, and one striped bass were collected.

Perch family (Percidae)

The Percidae family comprised 7 species, accounting for less than 0.1% of the total sample by weight and number. There were 6 darter species collected and the other species was sauger. Sauger ranged from 12.4 to 18.5 in TL and were collected at half of the sampling reaches. Sauger grew fast, reaching 18.5 in TL after three summers of growth (age 2).

Mooneye family (Hiodontidae)

Only one species of the Hiodontidae family was collected. Goldeye ranged from 11.9 to 17.2 in TL and were collected at 13 of 18 sampling reaches.

Sturgeon family (Acipenseridae)

Seventeen shovelnose sturgeon were collected at 8 of the 18 sampling reaches. Shovelnose sturgeon ranged from 8.3 to 29.8 in, measured from the tip of the snout to the fork in the caudal fin. No shovelnose were captured above RM 145.1.

Topminnow family (Fundulidae)

Blackstripe topminnow were the only members collected from the Fundulidae family. Blackstripe topminnow were collected at only one sampling reach (RM 45.7) and ranged from 1.4 to 2.2 in TL.

<u>Lamprey family (Petromyzontidae)</u>

The chestnut lamprey was the only member of this family collected, ranging from 7.4 to 11.1 in TL. Chestnut lamprey were collected at 4 of 18 sampling reaches, all above RM 73.6.

Bowfin family (Amiidae)

Two bowfin were collected at RM 45.7 and were 21.8 and 22.8 in TL.

Sculpin family (Cottidae)

Only one individual mottled sculpin was collected and it was from RM 141.5.

DISCUSSION

The overall purpose of this project was to collect fisheries and access data for the EFWR in 2003 (Hoffman 2006) and the WFWR in 2004. Major portions of the White River system had not been surveyed and other portions had been surveyed on an irregular basis. Data collected in this project would be used to develop a system-wide management plan. Another component of this project was to assess and recommend acquisition of new access sites in the area. There was one recommendation for acquiring a new access site at the end of CR 990 off HWY 231 in Owen County. A DFW crew was able to launch at an old ferry crossing and recommended the site for a boat ramp. Other sites that were on the acquisition list were in Greene, Owen, Monroe, and Morgan Counties.

There was no correlation between QHEI and IBI scores when all sample sites were pooled for the WFWR. For data from the EFWR 2003 (Hoffman 2006), there was a correlation between QHEI and IBI scores only when sample sites above Williams Dam were included (drainage area < 4,000 mi²). Drainage areas for the sample sites on the WFWR ranged from 2,700 to over 11,000 mi². There was no correlation between QHEI and IBI when the data for the WFWR was truncated for sites less than 4,000 mi². Similar to the EFWR, IBI scores for the WFWR were mostly considered "fair." The median IBI score for the WFWR was 40, which was slightly lower than the EFWR (median = 44). There was less variability in IBI scores for the WFWR (range = 32 to 46), compared to the EFWR, where IBI scores ranged from 30 to 58. For the EFWR and the WFWR, there was a significant, positive correlation between river mi and IBI. The median QHEI score for WFWR sample sites was 57, which was slightly lower than the median QHEI score (median = 64) for the EFWR. The highest QHEI score for the WFWR was 66 compared to 84 for the EFWR. The EFWR seems to be in better shape as far as fish diversity and habitat than the WFWR.

Combining electrofishing and seining yielded 70 species, representing 17 families of fish in the WFWR. In comparison, 69 species were collected in the EFWR in 2003 and 82 species were collected in the Wabash in 1999, although the samples for the Wabash were collected during July. No endangered species or species of special concern were collected in the WFWR.

No paddlefish were collected in the EFWR or the WFWR, but seven individual paddlefish were collected in the Wabash. The relative abundances of fish families represented in the total sample were expected. The sample consisted of mainly species from the Cyprinids in terms of abundance and Catostomids by weight, which is typical for large rivers. Crews captured 17 additional species via seines that were not captured with electrofishing. Six grass carp and one silver carp were collected via electrofishing. Grass carp were collected at 5 sampling reaches from RM 73 to 145. The one silver carp was collected at RM 15.9. Only one grass carp and no silver carp were collected in the EFWR.

Sportfish were collected in few numbers relative to other species, which was expected for larger rivers. Channel catfish, flathead catfish, and spotted bass were among the most abundant sportfish collected. Based on von Bertalannfy growth models, channel catfish grew slower in the WFWR (K = 0.138) compared to the EFWR (K = 0.165). Simulated results were similar for channel catfish in the WFWR compared to the EFWR and the Wabash. The current 10-in minimum size limit does not protect mature channel catfish. With the current estimates of annual mortality, exploitation was not large enough for growth overfishing to occur, but if exploitation increased by 10%, growth overfishing would likely occur. Flathead catfish in the EFWR and WFWR grew similarly, reaching almost 16 in TL by age 4. Freshwater drum were included in age and growth analyses because they were abundant in both the EFWR and the WFWR, and they received substantial directed effort during a creel survey on the EFWR (Hoffman 2004). Freshwater drum grew nearly identical in the EFWR and WFWR. Spotted bass growth was not compared between the EFWR and WFWR because a von Bertalannfy growth function could not be fit to data from the WFWR. However, spotted bass reached 11.5 in TL by age 4 in the WFWR compared to 10.7 in TL in the EFWR. Not enough information was collected to assess growth of other sport fish species, such as largemouth bass, white or black crappie, and smallmouth bass.

RECOMMENDATIONS

• Form a workgroup to review data collected from the EFWR and WFWR. The workgroup will develop a White River Basin management plan.

LITERATURE CITED

- Hoffman, K. J. 2006. White River Basin survey: East Fork White River, 2003. Indiana Department of Natural Resources, Division of Fish and Wildlife, Indianapolis, Indiana.
- Hoffman, K. J. 2004. Recreational use survey of the East Fork White and White Rivers, 2003. Indiana Department of Natural Resources, Division of Fish and Wildlife, Indianapolis, Indiana.
- Kingsley, D. W. 1983. Fisheries survey of the West Fork of White River and tributaries in Marion County. Indiana Department of Natural Resources, Division of Fish and Wildlife, Indianapolis, Indiana.
- Rankin, E. T. 1989. The Qualitative Habitat Evaluation Index (QHEI): Rationale, Methods and Application. Ohio Environmental Protection Agency, Columbus, Ohio, USA.
- Shipman, S., and seven coauthors. 2001. Manual of fisheries survey methods. Indiana Department of Natural Resources, Division of Fish and Wildlife, Indianapolis, Indiana.
- Simon, TP. 1998. Development of Index of Biotic Integrity expectations for the ecoregions of Indiana. IV. Eastern Corn Belt Plain. U.S. Environmental Protection Agency, Region V, Chicago, IL. EPA.

Submitted by: Kevin J. Hoffman, Asst. Research Biologist

Date: April 9, 2007

Approved by: Robert L. Ball, Research Biologist

Approved by: Non Schoenung, Fisheries Supervisor

Date: August 28, 2007

Table 1. Station water chemistry information, West Fork White River, September, 2004.

	Secchi	Air	Water	Dissolved	
River	disk	Temperature	Temperature	oxygen	Conductivity
Mile	(in)	(°F)	(°F)	(ppm)	(µS)
3.6	14	82.9	80.1	8.0	656
15.9	12	76.3	76.6	10.0	678
24.3	9	72.9	78.1	13.0	672
39.2	10	82.6	79.2	15.0	686
45.7	13	67.5	80.6	8.0	702
57.5	10	59.9	68.2	11.6	827
61.52	10	59.9	68.2	11.6	803
73.06	11	64.9	69.4	11.4	874
84.5	11	64.9	69.4	11.4	874
101.4	12	62.8	67.8	9.1	870
105.7	16	75.0	72.0	14.5	*
115.5	18	82.0	74.0	17.0	1,101
125.0	12	71.0	70.0	10.0	1,101
134.7	16	80.0	72.0	16.0	*
145.1	13	80.0	72.0	11.0	1,116
156.36	21	75.0	68.0	12.0	700
164.53	20	70.0	68.0	8 to 9	1,000
194.09	21	64.0	70.0	13 to 14	1,000

Table 2. Station location, drainage area, flow, gradient, sample distance, average width, average and maximum depth, West Fork White River, September, 2004.

River mile	County	Nearest town	Drainage area (mi2)	Flow (cfs)	Gradient (ft/mi)	Sample distance (ft)	Average width (ft)	Average depth (in)	Maximum depth (in)
3.6	Gibson/Pike	Mt. Carmel, IL	11,349	2580	0.4	2796	506.4	38.4	75.6
15.9	Gibson/Knox	Hazelton, IN	11,295	2200	0.4	4758	329.4	69.6	144.0
24.3	Gibson/Knox	Decker, IN	11,246	2250	0.4	5058	245.6	45.6	87.6
39.2	Knox/Pike	Petersburg, IN	11,170	2200	0.8	4740	444.0	54.0	118.8
45.7	Pike/Knox	Petersburg, IN	11,117	2150	0.9	1794	481.8	64.8	140.4
57.5	Daviess/Knox	Washington	5,254	3400	0.94	5050	318.6	36.4	81.6
61.52	Daviess/Knox	Washington	5,232	3400	1.02	3804	301.0	49.2	111.6
73.06	Daviess/Knox	Bicknell	5,032	3400	1.28	3511	217.5	55.3	264.0
84.5	Daviess/Knox	Edwardsport	4,852	3400	1.02	4803	289.0	45.3	100.8
101.4	Daviess/Knox	Elnora	4,793	3400	1.20	5870	223.5	33.7	102.0
105.7	Daviess/Knox/Greene	Elnora	4,769	908	0.93	4271	288.0	58.0	156.0
115.5	Greene	Plummer	4,658	908	0.93	3084	320.4	42.5	62.0
125.0	Greene	Bloomfield	4,413	908	0.93	4351	275.4	55.3	120.0
134.7	Greene	Worthington	3,184	908	1.45	2556	232.0	63.5	128.0
145.1	Owen	Farmers	3,091	837	1.14	3010	220.2	45.1	145.0+
156.36	Owen	Pottersville	2,999	*	1.20	3850	190.0	51.0	120.0
164.53	Owen	Spencer	2,968	*	1.23	3060	187.0	61.0	>132.0
194.09	Morgan	Martinsville	2,722	*	2.84	3060	234.0	41.0	96.0

Table 3. Station Qualitative Habitat Evaluation Index (QHEI) metric component scores, West Fork White River, September, 2004. Maximum scores are in parentheses.

River Mile	Substrate (20)	Cover (20)	Channel (20)	Riparian (10)	Pool (12)	Riffle (8)	Gradient (10)	Total (100)	Percent Pool	Percent Run	Percent Riffle
3.6	4	6	11	5	9	5	6	46.0	20	75	5
15.9	12	5	14	8.5	10	5	6	60.5	30	50	20
24.3	10	5	14	5	10	5	6	55.0	30	55	15
39.2	6	5	11	5.5	12	0	8	47.5	1	99	0
45.7	4	5	11	6	10	0	8	44.0	9	90	1
57.5	10	10	12	6	0	0	8	46.0	0	100	0
61.52	12	14	12	7	0	0	8	53.0	0	100	0
73.06	13	16	16	7	0	0	10	62.0	0	100	0
84.5	12	11	13	9	0	0	8	53.0	0	100	0
101.4	6	10	15	6	0	0	10	57.0	0	100	0
105.7	13.5	12	10	5	10	0	6	56.5	25	75	0
115.5	11.5	6	10	6	0	5	6	44.5	0	100	0
125.0	14	12	9	4.5	11	4	6	60.5	25	50	25
134.7	12	15	12	4.5	10	0	8	61.5	100	0	0
145.1	14	13	12	6	10	0	8	63.0	25	70	5
156.36	14	12	13	7	10	0	10	66.0	*	*	*
164.53	14	12	11	8	10	0	10	65.0	*	*	*
194.09	14	13	13	6	10	0	10	66.0	*	*	*

Table 4. Metric scores for the Index of Biologic Integrity (IBI) for each station, WFWR 2004.

		ı									4	River mile	9							
Triver Species 3	Metric	Metric description	3.6	15.9	24.3	39.2	45.7	57.5	61.52	73.06		101.4	105.7	5	125.0	134.7	145.1	156.36	164.53	194.09
Triver 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		Total # Species	3	κ	3	S	5	S	S	S	S	S	S	S	S	3	S	S	5	5
Total 5 1 1 3 5 3 3 3 3 3 3 3 3 4 <td></td> <td>% large river taxa</td> <td>5</td> <td>8</td> <td>5</td> <td>S</td> <td>5</td> <td>\$</td> <td>\$</td> <td>S</td> <td>\$</td> <td>\$</td> <td>S</td> <td>5</td> <td>5</td> <td>5</td> <td>5</td> <td>Ś</td> <td>S</td> <td>Ś</td>		% large river taxa	5	8	5	S	5	\$	\$	S	\$	\$	S	5	5	5	5	Ś	S	Ś
bodied vectors I I I I I I I I I I I I I I I I I I I		# of sunfish species	5	-	-	ω	5	ω	ω	8	ω	ω	\$	æ	æ		æ		ω	ю
Total 34 32 35 35 36 36 37 37 37 38 37 37 38 37 38 38 38 38 38 38 38 38 38 38 38 38 38		# round-bodied sucker species	-	-	1	-	-	-	ω	-	8	-	П	33	33	П	33	ю	ю	8
nt 1 2 3		# sensitive species	1	-	-	\mathcal{C}	3	κ	S	$^{\circ}$	т	κ		S	S	33	ю	S	8	2
vortes 3 4 3 4 3 3 4 3 4 3 4 3 4 <td></td> <td>% tolerant species</td> <td>1</td> <td>-</td> <td>-</td> <td>1</td> <td>-</td> <td>-</td> <td>-</td> <td>1</td> <td>1</td> <td>-</td> <td>-</td> <td>-</td> <td>-</td> <td>_</td> <td>-</td> <td>-</td> <td>_</td> <td>1</td>		% tolerant species	1	-	-	1	-	-	-	1	1	-	-	-	-	_	-	-	_	1
tivores 3 4 4 4 4 4 4 4 4 4 4 </td <td></td> <td>%omnivores</td> <td>3</td> <td>ϵ</td> <td>3</td> <td>\mathcal{E}</td> <td>3</td> <td>8</td> <td>ϵ</td> <td>3</td> <td>8</td> <td>ϵ</td> <td>-</td> <td>3</td> <td>3</td> <td>_</td> <td>3</td> <td>3</td> <td>3</td> <td>8</td>		%omnivores	3	ϵ	3	\mathcal{E}	3	8	ϵ	3	8	ϵ	-	3	3	_	3	3	3	8
vores 5 <td></td> <td>%insectivores</td> <td>3</td> <td>3</td> <td>1</td> <td>3</td> <td>3</td> <td>ϵ</td> <td>3</td> <td>3</td> <td>3</td> <td>3</td> <td>-</td> <td>3</td> <td>3</td> <td></td> <td>3</td> <td>8</td> <td>3</td> <td>3</td>		%insectivores	3	3	1	3	3	ϵ	3	3	3	3	-	3	3		3	8	3	3
He ils I I I I I I I I I I I I I I I I I I I		%carnivores	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
s 1 3 3 3 3 1 1 1 3 3 3 1 1 3 3 3 1 3		CPUE	-	-	8	ω	-	κ	\$	8	κ	\$	S	\$	5	5	5	\$	5	5
ss 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5		%simple lithophils	-	ω	ω	ω	-	-	ω	8		ω		ω	ω	ω	ω	8	ω	ϵ
34 32 32 40 38 38 46 40 40 42 36 46 46 34 44 44 46		%DELT anomalies	5	S	5	S	5	S	S	5	\$	S	ς.	ς.	\$	2	\$	\$	5	\$
		Total	34	32	32	40	38	38	46	40	40	42	36	46	46	34	44	44	46	46

Table 5. Number, mean length at age, and standard error determined from age-length keys for selected species in fall 2004, West Fork White River.

	, I						A	Age					
Common name	Statistic	-	2	3	4	\$	9	7	∞	6	10	11	12
Channel catfish	Mean SE N		11.4 0.36 10	13.9 0.42 10	15.8 0.54 7	17.2 0.21 25	19.6 0.47 15	21.6 0.66 7	21.5 1.08 5		20.5 NA 1		23.5 NA 1
Flathead catfish	Mean SE N	8.8 0.94 4	11.1 0.29 11	12.8 0.18 7	15.8 0.92 6	13.5 NA 1		18.8 0.75 4					
Freshwater drum	Mean SE N	8.9 0.20 9	12.1 0.19 60	14.3 0.14 96	16.7 0.15 10	16.2 0.19 6	18.0 0.29 4						
Spotted bass	Mean SE N	6.3 0.16 23	7.8 0.11 12	9.2 0.29 5	11.5 0.19 24			13.8 NA 1					




Figure 1. Sampling sites for the West Fork White River, 2004.




Figure 2. IBI scores and QHEI scores for each station (top) and IBI scores for each station based on river mile (bottom), WFWR 2004.




Figure 3. Length-frequency distribution of channel catfish, flathead catfish, freshwater drum, and spotted bass collected in the West Fork White River, 2004.

APPENDIX A

LIST OF ACCESS SITES ON THE WEST FORK WHITER RIVER

Daviess

NW of Elnora on CR 400E 5 mi W of Plainville on SR 358 to Carnahan Old US 50, W of Washington

Gibson

SR 56 at Main and First St. in Hazelton Mt. Carmel, IL launch site, NE on 4th St. from SR 1

Greene

1.1 mi SE of Worthington on SR 157

Hamilton

SR 19, 0.3 mi N of SR 38 at Forest Park in Noblesville 3.7 mi E of Strawtown 106th St., Hazel Dell Landing 116th St. Bridge River Bend Campground at Clare

Knox

NW of the SR 61 bridge in Petersburg East of Wheatland off of SR 150

Madison

Edgewater Park in Anderson at East 10th St. Mounds State Park, SR 232 E of Anderson

Marion

From 96th St to 16th St. with public access at city parks

Morgan

4.5 mi N of Martinsville on SR 37, then 0.7 mi W on Henderson 1 mi W of Waverly on SR 144 to Old SR 144

Owen

1.4 mi E of Spencer on SR 46, then 1 mi S on Old Southport Rd.

APPENDIX B

NAME, NUMBER, PERCENTAGE, SIZE, WEIGHT, AND OCCURRENCE INDEX OF FISHES COLLECTED, WEST FORK WHITE RIVER, FALL 2004

Appendix B. Name, number, percentage, size, weight, and occurrence index of fishes collected, West Fork White River, fall 2004 (all gears combined).

Common name	Scientific name	Total number	% by number	Min.	Max.	Total weight (lb)	% by weight	Occurrence index
Spotfin shiner	Cyprinella spiloptera	17763	66	0.6	3.1	23.6	0.6	18
Bullhead minnow	Pimephales vigilax	1961	7.3	0.6	3.1	3.03	0.1	16
Sand shiner	Notropis ludibundus	1512	5.6	0.8	2.6	1.84	0	15
Bluntnose minnow	Pimephales notatus	1004	3.7	0.7	2.9	1.7	0	18
Western mosquitofish	Gambusia affinis	781	2.9	0.5	1.9	0.99	0	15
River carpsucker	Carpiodes carpio	592	2.2	0.7	18.2	391.66	9.9	18
MS silvery minnow	Hybognathus nuchalis	460	1.7	1.8	4	3.85	0.1	15
Steelcolor shiner	Cyprinella whipplei	343	1.3	0.8	3.2	0.43	0	6
Brook silverside	Labidesthes sicculus	318	1.2	1.6	3	0.76	0	11
Gizzard shad	Dorosoma cepedianum	298	1.1	2.3	16.8	83.12	2.1	18
Smallmouth buffalo	Ictiobus bubalus	216	0.8	0.9	28.1	590.16	14.9	18
Freshwater drum	Aplodinotus grunniens	200	0.7	3.7	24.1	241	6.1	18
Common carp	Cyprinus carpio	182	0.7	13.9	28.7	1027.13	25.9	18
Silverjaw minnow	Notropis buccatus	182	0.7	1	2.6	0.35	0	9
Shortnose gar	Lepisosteus platostomus	181	0.7	16.9	29.5	226.6	5.7	17
Spotted bass	Micropterus punctulatus	101	0.4	2.2	14.2	29.64	0.7	16
Channel catfish	Ictalurus punctatus	95	0.4	1.8	25.6	154.19	3.9	16
Blue sucker	Cycleptus elongatus	77	0.3	9.8	30.7	350.26	8.8	14
Quillback	Carpiodes cyprinus	65	0.2	1.4	18.7	62.1	1.6	9
Longnose gar	Lepisosteus osseus	57	0.2	17.9	45	118.23	3	14
Flathead catfish	Pylodictis olivaris	46	0.2	4.8	33	67.75	1.7	15
Suckermouth minnow	Phenacobius mirabilis	45	0.2	2.5	3.6	0.6	0	6
Bluegill	Lepomis macrochirus	39	0.1	0.9	6	0.86	0	12
Bigmouth buffalo	Ictiobus cyprinellus	38	0.1	13.8	28.1	240.87	6.1	13
White bass	Morone chrysops	35	0.1	3.2	16.3	26	0.7	14
Goldeye	Hiodon alosoides	32	0.1	11.9	17.2	35.15	0.9	13
Longear sunfish	Lepomis megalotis	22	0.1	2	5.2	1.27	0	9
Emerald shiner	Notropis atherinoides	22	0.1	1.5	3.6	0.08	0	9
Central stoneroller	Campostoma pullum	22	0.1	2.3	3.3	0.19	0	6
Striped shiner	Luxilus chrysocephalus	20	0.1	2	2.6	0.09	0	3
Redfin shiner	Lythrurus umbratilis	19	0.1	1.1	2	0.01	0	5
Sauger	Sander canadense	18	0.1	12.4	18.5	19.87	0.5	9
Shovelnose sturgeon	Scaphirhynchus platorynchus	17	0.1	8.3	29.8	23.89	0.6	8
Golden redhorse	Moxostoma erythrurum	14	0.1	2.6	14.4	6.94	0.2	5
Shorthead redhorse	Moxostoma macrolepidotum	14	0.1	3.1	18.8	8.32	0.2	6
Black buffalo	Ictiobus niger	11	0	13.7	32.9	71.96	1.8	6
Dusky darter	Percina sciera	11	0	1.4	3	0.07	0	7
Highfin carpsucker	Carpiodes velifer	10	0	7.4	16.7	8.51	0.2	5
Blackstripe topminnow	Fundulus notatus	8	0	1.4	2.2	0.02	0	1
White crappie	Pomoxis annularis	6	0	6.1	12.8	2.4	0.1	5
Northern hogsucker	Hypentelium nigricans	6	0	4	5.4	0.21	0	1
Orangespotted sunfish	Lepomis humilis	6	0	2.3	3.9	0.22	0	3
Grass carp	Ctenopharyngodon idella	6	0	28	36.3	72.5	1.8	5
Chestnut lamprey	Ichthyomyzon castaneus	6	0	7.4	11.1	0.35	0	4
Yellow bass	Morone mississippiensis	6	0	3.4	9.1	0.47	0	6

Green sunfish	Lepomis cyanellus	5	0	1.4	2.4	0.02	0	3
Hybrid striped bass	Morone saxatilis x M. chrysops	5	0	16.1	26.5	19.42	0.5	3
Speckled chub	Macrhybopsis aestivalis	5	0	1.4	2.5	0.02	0	2
Silver redhorse	Moxostoma anisurum	4	0	13.7	24.1	15.23	0.4	4
Smallmouth bass	Micropterus dolomieu	4	0	7	12.5	2.07	0.1	2
Mountain madtom	Noturus eleutherus	4	0	1.6	2	0.02	0	2
Logperch	Percina caprodes	3	0	3.8	5.8	0.1	0	1
Slenderhead darter	Percina phoxocephala	3	0	2.4	3.2	0.02	0	1
Rosyface shiner	Notropis rubellus	3	0	1.5	1.8	*	0	1
Bowfin	Amia calva	2	0	21.8	22.8	7.3	0.2	1
Eastern sand darter	Ammocrypta pellucida	2	0	2	2	*	0	1
Bigeye chub	Notropis amblops	2	0	2.2	2.3	*	0	2
Hybrid sunfish	Lepomis macrochirus spp.	2	0	2.7	3.8	0.06	0	1
River shiner	Notropis blennius	1	0	2.7	2.7	0.01	0	1
Silver chub	Macrhybopsis storeriana	1	0	9.8	9.8	0.04	0	1
Largemouth bass	Micropterus salmoides	1	0	12.5	12.5	0.94	0	1
Black crappie	Pomoxis nigromaculatus	1	0	12.2	12.2	1.11	0	1
Black redhorse	Moxostoma duquesnei	1	0	3.2	3.2	0.01	0	1
Blue catfish	Ictalurus furcatus	1	0	20.2	20.2	2.5	0.1	1
Threadfin shad	Dorosoma petenense	1	0	4.1	4.1	0.02	0	1
Johnny darter	Etheostoma nigrum	1	0	1.7	1.7	0.01	0	1
Rainbow darter	Etheostoma caeruleum	1	0	1.7	1.7	0.01	0	1
Redear sunfish	Lepomis microlophus	1	0	2	2	*	0	1
Mottled sculpin	Cottus bairdi	1	0	1.6	1.6	0.01	0	1
Silver carp	Hypophthalmichthys molitrix	1	0	30.3	30.3	11	0.3	1
Striped bass	Morone saxatilis	1	0	29.5	29.5	10.75	0.3	1
Channel shiner	Notropis wickliffi	1	0	2.6	2.6	0.01	0	1
72 Species & 2 Hybrids		26925	100.0			3969.92	100.0	

*Less than 0.01 lbs.

APPENDIX C

SPECIES, NUMBER, AND WEIGHT OF FAMILIES COLLECTED FROM THE WEST FORK WHITE RIVER, FALL 2004.

Appendix C. Species, number, and weight of families collected from the West Fork White River, fall 2004. All gears combined.

2004. All ge	ears combined.				
Fam	ily	Number	%	Weight	%
Cyprinidae - carps and min Common carp Steelcolor shiner MS silvery minnow Spotfin shiner Emerald shiner River shiner Sand shiner Salver chub Central stoneroller Bluntnose minnow Suckermouth minnow	nows Bullhead minnow Grass carp Silver carp Striped shiner Channel shiner Redfin shiner Silverjaw minnow Rosyface shiner Speckled chub Bigeye chub	23555	87.5	1146.48	29.1
Catostomidae - Suckers Quillback Golden redhorse River carpsucker Shorthead redhorse Silver redhorse Blue sucker	Northern hogsucker Smallmouth buffalo Bigmouth buffalo Black buffalo Highfin carpsucker Black redhorse	1048	3.9	1746.23	43.9
Poecilidae - Livebearers Western mosquitofish		781	2.9	0.99	0.0
Atherinidae - Silversides Brook silverside		318	1.2	0.76	0.0
<u>Clupeidae - Herrings</u> Gizzard shad	Threadfin shad	299	1.1	83.14	2.1
<u>Lepisosteidae - Gars</u> Shortnose gar	Longnose gar	238	0.9	344.83	8.7
<u>Sciaenidae - Drums</u> Freshwater drum		200	0.7	241.00	6.1
Centrarchidae - Sunfishes Longear sunfish Bluegill White crappie Spotted bass Green sunfish Smallmouth bass	Largemouth bass Black crappie Orangespotted sunfish Redear sunfish Hybrid sunfish	188	0.7	38.59	1.0

Ictaluridae - Bullhead catfis Flathead catfish Channel catfish	<u>shes</u> Blue catfish Mountain madtom	146	0.5	224.46	5.6
Moronidae - Temperate ba White bass Hybrid striped bass	sses Yellow bass Striped bass	47	0.2	56.64	1.4
Percidae - Perches Logperch Slenderhead darter Sauger Dusky darter	Johnny darter Rainbow darter Eastern sand darter	39	0.1	20.08	0.5
<u>Hiodontidae - Mooneyes</u> Goldeye		32	0.1	35.15	0.9
Acipenseridae - Sturgeons Shovelnose sturgeon		17	0.1	23.89	0.6
<u>Fundulidae - Killifishes</u> Blackstripe topminnow		8	0.0	0.02	0.0
Petromyzontidae - Lampre Chestnut lamprey	<u>ys</u>	6	0.0	0.35	0.0
<u>Amiidae - Bowfins</u> Bowfin		2	0.0	7.3	0.2
Cottidae - Sculpins Mottled sculpin		1	0.0	0.01	0.0
Total		26,925	100.0	3,982.17	100.0

APPENDIX D

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Gizzard shad	23	16.2	2.3 - 15.7	12.12	6.1
Spotfin shiner	15	10.6	1.0 - 2.0	*	**
Freshwater drum	15	10.6	3.7 - 15.8	14.00	7.1
Shortnose gar	13	9.2	19.6 - 23.9	16.00	8.1
Common carp	12	8.5	22.3 - 27.0	79.57	40.1
Emerald shiner	10	7.0	1.5 - 2.1	*	**
Brook silverside	9	6.3	2.1 - 2.5	0.02	**
River carpsucker	9	6.3	12.8 - 18.5	16.17	8.1
Bluntnose minnow	7	4.9	0.8 - 2.8	0.02	**
Bigmouth buffalo	6	4.2	13.8 - 26.6	35.85	18.1
Spotted bass	5	3.5	2.9 - 9.7	0.75	0.4
Channel catfish	4	2.8	4.9 - 16.9	4.30	2.2
Western mosquitofish	4	2.8	0.8 - 1.1	*	**
Smallmouth buffalo	2	1.4	14.8 - 23.8	9.85	5.0
Blue sucker	1	0.7	23.8	4.48	2.3
Bluegill	1	0.7	1.5 - 4.3	0.14	0.1
Flathead catfish	1	0.7	11.2	0.52	0.3
Longear sunfish	1	0.7	2.3	0.01	**
Longnose gar	1	0.7	36.1	4.23	2.1
Redear sunfish	1	0.7	2.0	*	**
White bass	1	0.7	9.3	0.36	0.2
White crappie	1	0.7	6.1	0.08	**
Total - 22 Species	142	100.0		198.47	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	273	59.0	1.0 - 2.3	0.29	0.2
Mississippi silvery minnow	70	15.1	1.8 - 3.9	0.29	0.2
Blue sucker	17	3.7	9.8 - 27.5	69.60	37.0
Freshwater drum	15	3.2	9.5 - 16.7	15.67	8.3
Longnose gar	12	2.6	17.9 - 32.0	22.62	12.0
Shortnose gar	12	2.6	20.5 - 25.9	16.60	8.8
Bullhead minnow	8	1.7	1.0 - 2.1	0.02	**
White bass	8	1.7	9.7 - 16.1	7.91	4.2
Bluntnose minnow	6	1.3	1.6 - 2.0	0.01	**
Channel catfish	6	1.3	17.8 - 21.9	14.60	7.8
Gizzard shad	5	1.1	2.5 - 9.1	0.40	0.2
Sand shiner	5	1.1	0.8 - 1.8	*	**
River carpsucker	4	0.9	4.0 - 17.7	6.14	3.3
Suckermouth minnow	4	0.9	2.5 - 3.2	0.04	**
Western mosquitofish	4	0.9	0.8 - 1.7	*	**
Emerald shiner	3	0.6	2.8 - 3.1	0.01	**
Common carp	2	0.4	21.6 - 26.9	13.63	7.2
Shovelnose sturgeon	2	0.4	23.4 - 29.8	1.14	0.6
Silverjaw minnow	2	0.4	2.1 - 2.2	*	**
Smallmouth buffalo	2	0.4	12.4 - 22.0	6.99	3.7
Flathead catfish	1	0.2	9.6	0.30	0.2
Goldeye	1	0.2	13.4	0.77	0.4
Silver carp	1	0.2	30.3	11.00	5.9
Total - 23 Species	463	100.0		188.03	100.0

^{*}Less than 0.01 Lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Mississippi silvery minnow	180	33.9	1.8 - 3.3	1.39	1.0
River carpsucker	174	32.8	0.9 - 17.8	8.89	6.3
Bluntnose minnow	59	11.1	1.0 - 1.9	0.04	**
Spotfin shiner	39	7.3	0.8 - 2.4	0.02	**
Shortnose gar	11	2.1	17.0 - 23.3	12.66	8.9
Freshwater drum	9	1.7	10.0 - 15.5	10.55	7.4
Silverjaw minnow	9	1.7	1.2 - 2.3	0.02	**
Blue sucker	8	1.5	19.8 - 28.1	35.40	24.9
Longnose gar	6	1.1	29.8 - 31.8	16.59	11.7
Western mosquitofish	6	1.1	0.7 - 1.8	*	**
Gizzard shad	5	0.9	3.8 - 13.0	0.97	0.7
White bass	5	0.9	8.1 - 16.3	4.15	2.9
Channel catfish	4	0.8	17.6 - 20.6	9.45	6.7
Common carp	4	0.8	23.0 - 25.5	26.17	18.4
Goldeye	4	0.8	13.5 - 16.6	4.40	3.1
Shovelnose sturgeon	3	0.6	20.4 - 24.5	3.96	2.8
Emerald shiner	1	0.2	2.4	*	**
Flathead catfish	1	0.2	11	0.48	0.3
Sauger	1	0.2	17.4	1.34	0.9
Smallmouth buffalo	1	0.2	20.5	4.75	3.3
Spotted bass	1	0.2	11.8	0.86	0.6
Total - 21 Species	531	100.0		142.09	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	335	49.5	0.9 - 2.4	0.29	0.3
Bullhead minnow	128	18.9	0.9 - 2.3	0.19	0.2
River carpsucker	70	10.3	0.7 - 16.2	1.95	2.3
Mississippi silvery minnow	44	6.5	1.8 - 3.7	0.21	0.2
Freshwater drum	21	3.1	5.4 - 17.6	24.66	29.0
Bluntnose minnow	13	1.9	1.6 - 2.4	0.03	**
Western mosquitofish	10	1.5	0.8 - 1.5	0.01	**
Gizzard shad	9	1.3	3.2 - 13.9	2.60	3.1
Common carp	6	0.9	20.5 - 24.6	31.38	36.9
Flathead catfish	6	0.9	4.8 - 18.7	4.45	5.2
Silverjaw minnow	6	0.9	1.0 - 1.8	0.01	**
Shortnose gar	5	0.7	18.6 - 25.8	6.41	7.5
Longear sunfish	4	0.6	2.0 - 5.0	0.20	0.2
Spotted bass	4	0.6	3.5 - 7.9	0.46	0.5
Emerald shiner	3	0.4	2.7 - 3.3	0.02	**
Bluegill	2	0.3	2.3 - 2.5	0.02	**
Goldeye	2	0.3	11.9 - 15.0	1.47	1.7
Bigeye chub	1	0.1	2.3	*	**
Blue catfish	1	0.1	20.2	2.50	2.9
Blue sucker	1	0.1	20.9	2.91	3.4
Channel catfish	1	0.1	17.2	1.96	2.3
Dusky darter	1	0.1	2.3	0.01	**
Orangespotted sunfish	1	0.1	2.3	0.01	**
Sand shiner	1	0.1	1.7	*	**
Smallmouth buffalo	1	0.1	17.2	2.50	2.9
White bass	1	0.1	12.3	0.81	1.0
Total - 26 Species	677	100.0		85.06	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Western mosquitofish	132	42.9	0.7 - 1.4	0.07	**
Spotfin shiner	52	16.9	0.9 - 2.5	0.07	**
Brook silverside	20	6.5	1.6 - 2.7	0.04	**
Freshwater drum	16	5.2	6.2 - 17.8	20.97	14.2
Gizzard shad	11	3.6	3.4 - 16.8	4.09	2.8
Common carp	10	3.2	19.9 - 22.6	42.52	28.7
Blackstripe topminnow	8	2.6	1.4 - 2.2	0.02	**
Flathead catfish	7	2.3	7.5 - 17.3	4.09	2.8
River carpsucker	7	2.3	15.1 - 17.8	14.28	9.6
Shortnose gar	7	2.3	18.0 - 24.4	8.34	5.6
Smallmouth buffalo	6	1.9	12.6 - 22.4	22.32	15.1
Spotted bass	5	1.6	2.9 - 12.0	1.52	1.0
Bluegill	3	1.0	0.9 - 1.5	*	**
Bluntnose minnow	3	1.0	0.7 - 1.6	*	**
Bullhead minnow	3	1.0	1.0 - 1.9	*	**
Longear sunfish	3	1.0	4.0 - 4.9	0.23	0.2
Longnose gar	3	1.0	27.5 - 45.0	16.55	11.2
Orangespotted sunfish	3	1.0	2.3 - 2.9	0.01	**
Bowfin	2	0.6	21.8 - 22.8	7.30	4.9
Eastern sand darter	2	0.6	2.0	*	**
Black crappie	1	0.3	12.2	1.11	0.7
Goldeye	1	0.3	14.4	0.88	0.6
Hybrid striped bass	1	0.3	16.1	1.67	1.1
Largemouth bass	1	0.3	12.5	0.94	0.6
White crappie	1	0.3	12.8	1.14	0.8
Total - 25 Species	308	100.0		148.16	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

NAME OF STREAM: WFWR

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Shortnose gar	31	32.6	19.0 - 28.3	38.28	20.8
River carpsucker	17	17.9	2.6 - 17.8	19.68	10.7
Freshwater drum	8	8.4	10.1 - 15.5	7.61	4.1
Smallmouth buffalo	7	7.4	12.5 - 24.0	27.64	15.0
Common carp	5	5.3	18.1 - 23.5	22.56	12.3
Channel catfish	5	5.3	15.4 - 23.0	11.12	6.0
Black buffalo	4	4.2	20.4 - 32.9	38.00	20.7
Longnose gar	3	3.2	25.7 - 26.5	4.15	2.3
Spotted bass	2	2.1	6.4 - 8.2	0.42	0.2
Spotfin shiner	2	2.1	2.1 - 2.3	0.02	*
Bullhead minnow	2	2.1	1.9 - 2.9	0.02	*
Blue sucker	1	1.1	26.6	5.75	3.1
Quillback	1	1.1	18.7	3.45	1.9
Bigmouth buffalo	1	1.1	17.7	3.21	1.7
Goldeye	1	1.1	16.8	1.30	0.7
Flathead catfish	1	1.1	12.4	0.68	0.4
Shovelnose sturgeon	1	1.1	8.3	0.05	*
Bluegill	1	1.1	3.0	0.02	*
Threadfin shad	1	1.1	4.1	0.02	*
Yellow bass	1	1.1	3.4	0.01	*
Totals - 20 Species	95			183.99	

^{*}Less than 0.1%.

DATE:9/21/04 STATION: RM 57.5 seine

NAME OF STREAM: WFWR

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	475	70.7	0.7 - 2.3	0.55	47.0
Bullhead minnow	73	10.9	0.9 - 2.6	0.13	11.1
Mississippi silvery minnow	50	7.4	2.2 - 3.4	0.34	29.1
Sand shiner	42	6.3	1.1 - 1.9	0.07	6.0
Western mosquitofish	19	2.8	0.8 - 1.2	0.01	0.9
Bluntnose minnow	5	0.7	1.6 - 1.9	0.01	0.9
Suckermouth minnow	3	0.4	2.5 - 2.7	0.02	1.7
Channel catfish	2	0.3	1.9 - 2.0	0.01	0.9
River carpsucker	1	0.1	2.1	0.01	0.9
Emerald shiner	1	0.1	2.4	0.01	0.9
Speckled chub	1	0.1	1.4	0.01	0.9
Totals - 11 Species	672		1	1.17	0.5

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Freshwater drum	20	14.8	5.4 - 17.9	21.30	8.8
Common carp	16	11.9	15.3 - 27.2	77.87	32.3
Smallmouth buffalo	13	9.6	12.3 - 23.4	50.78	21.0
Shortnose gar	11	8.1	16.9 - 27.1	14.50	6.0
Spotted bass	10	7.4	3.1 - 9.9	1.50	0.6
Gizzard shad	9	6.7	3.2 - 12.3	2.67	1.1
River carpsucker	8	5.9	4.3 - 17.3	7.39	3.1
Channel catfish	7	5.2	3.5 - 21.9	8.22	3.4
Bigmouth buffalo	5	3.7	19.9 - 24.5	32.50	13.5
Bluegill	5	3.7	1.9 - 4.1	0.13	0.1
Shovelnose sturgeon	4	3.0	6.9 - 26.7	5.02	2.1
Bullhead minnow	4	3.0	2.2 - 2.9	0.03	*
Flathead catfish	3	2.2	12.8 - 19.3	4.50	1.9
Goldeye	3	2.2	15.6 - 17.1	3.88	1.6
Longear sunfish	3	2.2	2.2 - 3.8	0.10	*
Mississippi silvery minnow	3	2.2	3.4 - 3.7	0.07	*
Longnose gar	2	1.5	21.2 - 34.3	3.83	1.6
Spotfin shiner	2	1.5	2.2 - 2.4	0.02	*
Blue sucker	1	0.7	23.2	3.50	1.4
Quillback	1	0.7	18.1	2.92	1.2
Highfin carpsucker	1	0.7	11.2	0.61	0.3
Silver chub	1	0.7	4.8	0.04	*
Yellow bass	1	0.7	3.6	0.01	*
Green sunfish	1	0.7	2.3	0.01	*
Emerald shiner	1	0.7	3.6	0.01	*
Totals - 25 Species	135			241.41	

^{*}Less than 0.1%.

DATE:9/21/04 STATION: RM 61.52 seine

NAME OF STREAM: WFWR

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	663	85.2	0.6 - 2.8	1.79	80.6
Bullhead minnow	32	4.1	1.2 - 2.5	0.07	3.2
Mississippi silvery minnow	23	3.0	2.2 - 3.1	0.13	5.9
Western mosquitofish	14	1.8	0.4 - 1.0	0.01	0.5
Sand shiner	12	1.5	1.5 - 2.0	0.02	0.9
Brook silverside	9	1.2	2.0 - 2.4	0.02	0.9
Suckermouth minnow	8	1.0	2.7 - 3.2	0.06	2.7
Speckled chub	4	0.5	1.7 - 2.5	0.01	0.5
Bluntnose minnow	4	0.5	1.0 - 2.2	0.01	0.5
Shorthead redhorse	3	0.4	3.4 - 3.8	0.05	2.3
Golden redhorse	3	0.4	2.6 - 3.2	0.02	0.9
Steelcolor shiner	1	0.1	1.7	0.01	0.5
Channel shiner	1	0.1	2.6	0.01	0.5
Black redhorse	1	0.1	3.2	0.01	0.5
Totals - 14 Species	778			2.22	

NAME, NUMBER, PERCENTAGE, SIZE, AND WEIGHT OF FISHES COLLECTED

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Mississippi silvery minnow	19	13.0	2.8 - 3.7	0.31	0.1
Common carp	14	9.6	16.5 - 26.5	70.30	27.9
Smallmouth buffalo	12	8.2	12.5 - 22.8	31.87	12.7
Shortnose gar	12	8.2	17.5 - 23.2	13.12	5.2
Freshwater drum	11	7.5	4.9 - 18.9	16.33	6.5
Gizzard shad	9	6.2	3.8 - 11.1	1.06	0.4
Blue sucker	8	5.5	15.8 - 26.8	25.87	10.3
Channel catfish	8	5.5	16.8 - 23.4	19.37	7.7
River carpsucker	8	5.5	8.1 - 18.1	14.72	5.8
Spotted bass	8	5.5	5.6 - 12.3	2.46	1.0
Bullhead minnow	8	5.5	2.0 - 2.9	0.05	*
Quillback	6	4.1	9.2 - 18.0	10.28	4.1
Flathead catfish	5	3.4	9.2 - 18.6	6.20	2.5
White bass	3	2.1	4.5 - 13.1	1.20	0.5
Bigmouth buffalo	2	1.4	23.3 - 27.1	21.00	8.3
Shovelnose sturgeon	2	1.4	24.3 - 25.7	4.38	1.7
Goldeye	2	1.4	15.3 - 16.4	2.67	1.1
Spotfin shiner	2	1.4	2.2 - 2.4	0.02	*
Grass carp	1	0.7	28.0	9.50	3.8
Sauger	1	0.7	15.1	1.04	0.4
Chestnut lamprey	1	0.7	7.9	0.04	*
Bluegill	1	0.7	4.0	0.03	*
Emerald shiner	1	0.7	3.4	0.02	*
Sand shiner	1	0.7	2.1	0.01	*
River shiner	1	0.7	2.7	0.01	*
Totals - 25 Species	146			251.86	

Less than 0.1%.

DATE:9/22/04 STATION: RM 73.06 seine

NAME OF STREAM: WFWR

CONTROLLANT) H D (DED	DED CENTER CE	SIZE RANGE	TOTAL WEIGHT	DED CENTA CE
COMMON NAME	NUMBER	PERCENTAGE	(INCHES)	(POUNDS)	PERCENTAGE
Bullhead minnow	140	48.1	0.9 - 2.6	0.29	43.9
Western mosquitofish	74	25.4	0.9 - 1.8	0.07	10.6
Brook silverside	47	16.2	2.0 - 2.6	0.09	13.6
Spotfin shiner	16	5.5	1.0 - 2.2	0.03	4.5
Sand shiner	8	2.7	1.2 - 2.1	0.02	3.0
Gizzard shad	3	1.0	4.3 - 4.4	0.09	13.6
Bluntnose minnow	2	0.7	2.1 - 2.3	0.01	1.5
Freshwater drum	1	0.3	4.9	0.06	9.1
Totals - 8 Species	291			0.66	

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Gizzard shad	28	18.1	4.0 - 13.5	5.40	1.7
Common carp	19	12.3	15.9 - 25.3	91.48	29.0
Freshwater drum	14	9.0	10.7 - 17.6	18.01	5.7
River carpsucker	11	7.1	9.0 - 18.2	13.14	4.2
Channel catfish	10	6.5	10.4 - 24.9	17.99	5.7
Blue sucker	9	5.8	22.6 - 27.6	48.25	15.3
Smallmouth buffalo	9	5.8	18.6 - 20.9	36.10	11.5
Mississippi silvery minnow	9	5.8	3.0 - 3.7	0.21	0.1
Spotted bass	7	4.5	6.0 - 10.2	1.68	0.5
Goldeye	6	3.9	13.5 - 16.1	5.98	1.9
Flathead catfish	6	3.9	6.6 - 13.6	3.68	1.2
Bigmouth buffalo	4	2.6	20.0 - 28.1	28.50	9.0
Quillback	4	2.6	9.8 - 16.8	7.49	2.4
Shortnose gar	4	2.6	18.5 - 24.8	6.46	2.0
Longnose gar	4	2.6	20.6 - 26.8	4.86	1.5
Bullhead minnow	4	2.6	2.1 - 3.1	0.02	*
Grass carp	1	0.6	36.3	19.50	6.2
Black buffalo	1	0.6	21.1	5.00	1.6
White bass	1	0.6	12.5	0.86	0.3
Shorthead redhorse	1	0.6	9.9	0.30	0.1
White crappie	1	0.6	8.0	0.24	0.1
Chestnut lamprey	1	0.6	7.4	0.03	*
Bluegill	1	0.6	3.2	0.02	*
Totals - 23 Species	155			315.20	

^{*}Less than 0.1%.

DATE:9/20/04 STATION: RM 84.50 seine

NAME OF STREAM: WFWR

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	445	89.9	0.9 - 2.8	0.72	78.3
Bullhead minnow	21	4.2	0.9 - 2.4	0.06	6.5
Bluntnose minnow	6	1.2	1.4 - 2.1	0.02	2.2
Sand shiner	6	1.2	1.3 - 1.9	0.01	1.1
Brook silverside	5	1.0	2.0 - 2.6	0.01	1.1
Mississippi silvery minnow	3	0.6	3.2 - 3.5	0.04	4.3
Redfin shiner	3	0.6	1.3 - 1.5	0.01	1.1
Steelcolor shiner	2	0.4	1.2 - 1.3	0.01	1.1
River carpsucker	1	0.2	2.2	0.01	1.1
Channel catfish	1	0.2	1.8	0.01	1.1
Johnny darter	1	0.2	1.7	0.01	1.1
Suckermouth minnow	1	0.2	2.8	0.01	1.1
Totals - 12 Species	495			0.92	

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Mississippi silvery minnow	32	18.6	2.9 - 3.8	0.48	0.2
River carpsucker	31	18.0	1.4 - 16.6	22.90	10.2
Gizzard shad	18	10.5	4.2 - 13.9	3.72	1.7
Blue sucker	17	9.9	20.1 - 30.7	86.57	38.7
Freshwater drum	9	5.2	8.9 - 16.4	10.61	4.7
Shortnose gar	8	4.7	20.0 - 23.2	8.99	4.0
Spotfin shiner	8	4.7	1.5 - 2.5	0.02	*
Smallmouth buffalo	7	4.1	15.0 - 23.1	25.20	11.3
Quillback	6	3.5	8.7 - 16.4	7.93	3.5
Longnose gar	6	3.5	20.7 - 26.1	6.15	2.7
Common carp	5	2.9	19.1 - 23.1	24.44	10.9
Sand shiner	4	2.3	1.4 - 1.9	0.01	*
Bullhead minnow	3	1.7	2.6 - 2.7	0.02	*
Channel catfish	2	1.2	19.4 - 21.3	4.76	2.1
Shovelnose sturgeon	2	1.2	24.8 - 25.8	4.63	2.1
Goldeye	2	1.2	16.2 - 16.7	2.65	1.2
Sauger	2	1.2	14.7 - 15.5	1.81	0.8
White bass	2	1.2	9.4 - 12.0	1.11	0.5
Highfin carpsucker	2	1.2	7.4 - 7.6	0.42	0.2
Bigmouth buffalo	1	0.6	23.5	9.25	4.1
Black buffalo	1	0.6	15.4	1.83	0.8
Spotted bass	1	0.6	6.4	0.14	0.1
Longear sunfish	1	0.6	4.9	0.09	*
Yellow bass	1	0.6	4.1	0.03	*
Mountain madtom	1	0.6	1.7	0.01	*
Totals - 25 Species *Less than 0.1%	172			223.77	

^{*}Less than 0.1%.

DATE:9/20/04 STATION: RM 101.4 seine

NAME OF STREAM: WFWR

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
			(, , , , , , , , , , , , , , , , , , ,	(
Spotfin shiner	725	73.8	0.6 - 2.7	0.80	57.7
Sand shiner	187	19.0	0.8 - 2.1	0.32	23.1
Bullhead minnow	55	5.6	0.6 - 1.3	0.02	1.4
Mississippi silvery minnow	7	0.7	3.4 - 3.5	0.11	7.9
River carpsucker	4	0.4	1.7 - 4.2	0.09	6.5
Silverjaw minnow	1	0.1	2.1	0.01	0.7
Bluntnose minnow	1	0.1	1.3	0.01	0.7
Stoneroller	1	0.1	2.3	0.01	0.7
Western mosquitofish	1	0.1	1.0	0.01	0.7
Emerald shiner	1	0.1	3.3	0.01	0.5
Totals - 10 Species	983			1.39	

STATION: RM 105.7, Elnora electrofishing

NAME OF STREAM: WFWR

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Common carp	19	14.4	13.9 - 26.0	104.12	43.1
Quillback	17	12.9	8.4 - 16.2	21.96	9.1
Smallmouth buffalo	14	10.6	11.0 - 27.0	35.67	14.8
Gizzard shad	13	9.8	4.4 - 10.7	2.39	1.0
Shortnose gar	11	8.3	18.6 - 23.0	12.00	5.0
River carpsucker	9	6.8	1.3 - NA	12.50	5.2
Freshwater drum	6	4.5	5.7 - 17.4	9.20	3.8
Spotted bass	6	4.5	2.2 - 10.4	0.83	0.3
Channel catfish	5	3.8	7.9 - 21.6	8.73	3.6
Bluegill	5	3.8	2.1 - 6.0	0.22	0.1
Sand shiner	3	2.3	2.2	*	**
Longnose gar	3	2.3	19.9 - 26.3	2.75	1.1
Flathead catfish	3	2.3	11.4 - 21.1	4.68	1.9
White bass	3	2.3	3.2 - 12.9	1.04	0.4
Grass carp	2	1.5	29.3 - 34.0	22.50	9.3
Goldeye	2	1.5	14.2 - 15.3	1.96	0.8
Orangespotted sunfish	2	1.5	2.5	0.20	0.1
Mississippi silvery minnow	3	1.5	3.6	0.02	**
Green sunfish	2	1.5	2.1 - 2.4	0.01	**
Central stoneroller	1	0.8	2.8	*	**
Bluntnose minnow	1	0.8	1.8	*	**
Hybrid sunfish	1	0.8	3.8	0.05	**
White crappie	1	0.8	10.6	0.57	0.2
Total - 22 Species and 1 hybrid	132	100.0		241.40	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	1158	76.1	1.0 - 2.5	0.95	59.0
Sand shiner	207	13.6	1.2 - 2.4	0.32	19.9
Bluntnose minnow	91	6.0	0.9 - 2.6	0.02	1.2
Bullhead minnow	52	3.4	1.3 - 3.0	0.03	1.9
Dusky darter	5	0.3	1.4 - 2.8	0.05	3.1
Quillback	4	0.3	1.8 - 2.4	0.20	12.4
River carpsucker	4	0.3	1.3 - 2.8	0.01	0.6
Gizzard shad	1	0.1	4.4	0.03	1.9
Total - 8 Species	1522	100.0		1.61	100.0

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
River carpsucker	31	24.6	11.0 - 17.4	52.10	19.2
Common carp	19	15.1	18.0 - 28.5	118.75	43.7
Freshwater drum	17	13.5	10.4 - 24.1	29.71	10.9
Spotted bass	12	9.5	3.2 - 12.0	3.40	1.3
Channel catfish	8	6.3	10.4 - 22.0	12.47	4.6
Longear sunfish	6	4.8	3.5 - 5.2	0.44	0.2
Gizzard shad	6	4.8	10.0 - 13.0	2.75	1.0
Smallmouth buffalo	3	2.4	17.5 - 23.2	14.94	5.5
Bluntnose minnow	3	2.4	1.4 - 2.7	*	**
Slenderhead darter	3	2.4	2.4 - 3.2	0.02	**
Flathead catfish	2	1.6	14.5 - 16.0	2.63	1.0
Bigmouth buffalo	2	1.6	18.3 - 24.3	10.70	3.9
Chestnut lamprey	2	1.6	7.9 - 11.1	0.15	0.1
Black buffalo	1	0.8	24.3	5.75	2.1
White bass	1	0.8	10.3	0.47	0.2
Sauger	1	0.8	15.5	1.10	0.4
Highfin carpsucker	1	0.8	13.0	1.15	0.4
Mirror carp	1	0.8	29.0	12.25	4.5
Silver redhorse	1	0.8	13.7	1.05	0.4
Shorthead redhorse	1	0.8	17.8	1.77	0.7
Hybrid sunfish	1	0.8	2.7	0.01	**
Bluegill	1	0.8	2.1	*	**
Spotfin shiner	1	0.8	2.2	*	**
Mississippi silvery minnow	1	0.8	3.7	0.02	**
Dusky darter	1	0.8	1.8	*	**
Total - 24 Species and 1 hybrid	126	100.0		271.63	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

STATION: RM 115.5, Plummer seine

NAME OF STREAM: WFWR

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
			(0.00000)	(2 2 2 1 2 2)	
Spotfin shiner	386	39.3	0.7 - 2.2	0.23	13.6
Western mosquitofish	292	29.8	0.5 - 1.7	0.12	7.1
Bullhead minnow	247	25.2	0.9 - 1.3	0.15	8.9
Sand shiner	19	1.9	0.9 - 2.6	*	**
Brook silverside	13	1.3	2.1 - 3.0	0.04	2.4
Bluntnose minnow	11	1.1	1.0 - 2.4	0.06	3.6
Redfin shiner	11	1.1	1.2 - 2.0	*	**
Gizzard shad	1	0.1	3.2	0.02	1.2
Freshwater drum	1	0.1	13.4	1.07	63.3
Total - 9 Species	981	100.0		1.69	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Shortnose gar	17	14.0	21.8 - 29.5	24.00	8.3
Common carp	16	13.2	19.9 - 28.7	89.50	30.9
River carpsucker	15	12.4	10.1 - 16.7	23.00	7.9
Smallmouth buffalo	14	11.6	13.5 - 28.0	65.50	22.6
Gizzard shad	14	11.6	9.8 - 12.1	5.50	1.9
Freshwater drum	9	7.4	9.2 - 15.3	8.35	2.9
Longnose gar	6	5.0	25.2 - 30.5	9.20	3.2
Spotted bass	4	3.3	3.7 - 14.2	3.13	1.1
Blue sucker	4	3.3	22.9 - 29.8	23.82	8.2
Channel catfish	3	2.5	12.7 - 22.6	3.98	1.4
White bass	3	2.5	12.0 - 15.3	3.36	1.2
Hybrid striped bass	3	2.5	16.8 - 26.5	10.75	3.7
White crappie	2	1.7	7.1 - 8.7	0.37	0.1
Sauger	2	1.7	14.2 - 18.4	2.19	0.8
Golden redhorse	2	1.7	13.2 - 13.7	1.93	0.7
Shorthead redhorse	2	1.7	15.7 - 17.3	3.45	1.2
Silver redhorse	1	0.8	24.1	5.50	1.9
Bigmouth buffalo	1	0.8	19.9	4.20	1.4
Longear sunfish	1	0.8	4.8	0.05	*
Shovelnose sturgeon	1	0.8	24.9	1.85	0.6
Yellow bass	1	0.8	9.1	0.36	0.1
Total - 20 Species and 1 hybrid	121	100.0		289.99	100.0

^{*}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	4614	83.8	0.8 - 2.8	7.82	75.8
Sand shiner	368	6.7	1.1 -2.4	0.77	7.5
Bullhead minnow	182	3.3	1.7 - 2.6	0.33	3.2
Brook silverside	132	2.4	1.7 - 2.8	0.40	3.9
Western mosquitofish	104	1.9	0.6 - 1.8	0.61	5.9
Bluntnose minnow	39	0.7	1.2 - 2.7	0.18	1.7
River carpsucker	29	0.5	1.3 - 2.7	0.07	0.7
Striped shiner	14	0.3	2.5 - 2.6	0.06	0.6
Mississippi silvery minnow	4	0.1	3.2 - 3.5	0.03	0.3
Steelcolor shiner	4	0.1	1.2 - 2.4	0.01	0.1
Central stoneroller	3	0.1	2.8 - 3.2	*	**
Silverjaw minnow	3	0.1	2.3 - 2.6	*	**
Mountain madtom	3	0.1	1.6 - 2.0	0.01	0.1
Shorthead redhorse	2	0.0	3.1 - 3.2	0.02	0.2
Green sunfish	2	0.0	1.4 - 1.7	*	**
Gizzard shad	1	0.0	2.3	0.01	0.1
Dusky darter	1	0.0	1.4	*	**
Redfin shiner	1	0.0	1.1	*	**
Emerald shiner	1	0.0	2.1	*	**
Total - 19 Species	5507	100.0		10.32	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	120	61.9	1.3 - 2.4	0.35	0.2
Gizzard shad	14	7.2	8.9 - 12.2	5.68	2.6
Smallmouth buffalo	12	6.2	11.6 - 28.1	73.50	33.1
River carpsucker	9	4.6	13.9 - 18.1	15.50	7.0
Shortnose gar	8	4.1	16.9 - 22.6	9.50	4.3
Common carp	6	3.1	17.7 - 26.7	37.48	16.9
Longnose gar	5	2.6	23.1 - 37.0	9.80	4.4
Goldeye	5	2.6	14.4 - 17.2	5.92	2.7
Bigmouth buffalo	5	2.6	23.7 - 24.6	32.25	14.5
Freshwater drum	3	1.5	13.1 - 15.7	3.84	1.7
Blue sucker	2	1.0	21.3 - 24.7	8.13	3.7
Grass carp	1	0.5	31.0	10.00	4.5
Hybrid striped bass	1	0.5	25.5	7.00	3.2
Yellow bass	1	0.5	3.8	0.03	*
Sauger	1	0.5	17.8	1.42	0.6
White bass	1	0.5	15.9	1.42	0.6
Total -15 Species and 1 hybrid	194	100.0		221.82	100.0

^{*}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
			,		
Spotfin shiner	938	79.0	1.1 - 2.4	0.72	74.2
Bullhead minnow	139	11.7	1.0 - 1.8	0.07	7.2
Brook silverside	29	2.4	2.1 - 2.7	*	**
Sand shiner	28	2.4	0.8 - 1.8	0.01	1.0
Western mosquitofish	19	1.6	0.7 - 1.4	0.01	1.0
Bluntnose minnow	16	1.3	0.9 - 2.2	0.01	1.0
Mississippi silvery minnow	8	0.7	3.0 - 3.8	0.12	12.4
River carpsucker	6	0.5	1.5 - 1.9	0.01	1.0
Smallmouth buffalo	3	0.3	1.0 - 1.1	*	**
Gizzard shad	1	0.1	4.3	0.02	2.1
Total - 10 Species	1187	100.0		0.97	100.0

^{*}Less than 0.01 lb.
**Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
River carpsucker	28	28.6	10.6 - 17.2	43.50	20.5
Shortnose gar	12	12.2	19.5 - 24.0	14.92	7.0
Flathead catfish	6	6.1	11.9 - 33.0	19.59	9.3
Freshwater drum	6	6.1	13.5 - 21.0	11.82	5.6
Gizzard shad	6	6.1	8.1 - 11.3	2.08	1.0
Common carp	6	6.1	22.4 - 26.4	44.00	20.8
Spotted bass	5	5.1	3.4 - 10.4	0.74	0.3
Blue sucker	5	5.1	23.5 - 27.6	20.43	9.6
Highfin carpsucker	5	5.1	11.0 - 16.7	5.87	2.8
Channel catfish	3	3.1	11.0 - 19.9	4.49	2.1
Bigmouth buffalo	3	3.1	19.0 - 22.9	17.08	8.1
Longnose gar	3	3.1	24.4 - 26.1	3.85	1.8
Goldeye	2	2.0	14.9	2.26	1.1
Shovelnose sturgeon	2	2.0	24.4 - 28.6	2.86	1.4
Bluegill	1	1.0	6.0	0.21	0.1
Grass carp	1	1.0	33.0	11.00	5.2
Silver redhorse	1	1.0	20.2	3.53	1.7
Smallmouth buffalo	1	1.0	21.9	3.50	1.7
Bluntnose minnow	1	1.0	2.1	*	**
Spotfin shiner	1	1.0	2.5	*	**
Total - 20 Species	98	100.0		211.73	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	556	63.0	0.8 - 2.9	0.56	46.3
Bluntnose minnow	105	11.9	1.0 - 2.9	0.20	16.5
Sand shiner	97	11.0	1.2 - 2.1	0.13	10.7
Bullhead minnow	69	7.8	1.0 - 1.6	0.05	4.1
Western mosquitofish	27	3.1	1.0 - 1.9	0.02	1.7
River carpsucker	10	1.1	1.8 - 3.2	0.11	9.1
Spotted bass	4	0.5	2.9 - 4.2	0.06	5.0
Silverjaw minnow	3	0.3	1.2 - 1.4	*	**
Redfin shiner	3	0.3	1.1 - 1.2	*	**
Yellow bass	1	0.1	4.4	0.03	2.5
Dusky darter	1	0.1	2.2	*	**
Gizzard shad	1	0.1	5.0	0.04	3.3
Bluegill	1	0.1	2.2	*	**
Mottled sculpin	1	0.1	1.6	*	**
Brook silverside	1	0.1	2.6	*	**
Mississippi silvery minnow	1	0.1	3.0	0.01	0.8
Smallmouth buffalo	1	0.1	0.9	*	**
Total -17 Species	882	100.0		1.21	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

COMMONINAME	NUMBER	DED CENTA CE	SIZE RANGE	TOTAL WEIGHT	DED CENTA CE
COMMON NAME	NUMBER	PERCENTAGE	(INCHES)	(POUNDS)	PERCENTAGE
River carpsucker	46	31.1	10.7 - 17.6	69.78	21.8
Gizzard shad	34	23.0	4.5 - 14.1	13.99	4.4
Smallmouth buffalo	17	11.5	13.5 - 25.3	92.36	28.9
Freshwater drum	11	7.4	5.5 - 14.5	8.75	2.7
Common carp	9	6.1	19.9 - 26.7	65.50	20.5
Shortnose gar	8	5.4	20.3 - 22.0	8.89	2.8
Bigmouth buffalo	3	2.0	21.7 - 22.7	20.44	6.4
Spotted bass	3	2.0	4.1 - 13.7	2.37	0.7
Blue sucker	2	1.4	26.0	12.63	3.9
Flathead catfish	2	1.4	11.0 - 19.9	3.30	1.0
Sauger	2	1.4	13.8 - 17.7	2.15	0.7
Channel catfish	2	1.4	10.3 - 14.3	1.09	0.3
White bass	2	1.4	8.8 - 10.0	0.69	0.2
Black buffalo	1	0.7	27.2	10.75	3.4
Longnose gar	1	0.7	31.8	3.15	1.0
Quillback	1	0.7	14.7	1.41	0.4
Golden redhorse	1	0.7	13.3	1.06	0.3
Goldeye	1	0.7	14.8	1.01	0.3
Highfin carpsucker	1	0.7	10.0	0.46	0.1
Dusky darter	1	0.7	3.0	0.01	*
Total - 20 Species	148	100.0		319.79	100.0

^{*}Less than 0.1%.

NAME OF STREAM: West Fork White River

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	2,906	74.0	0.6 - 2.9	3.64	62.1
Bullhead minnow	488	12.4	0.8 - 3.0	1.14	19.5
Bluntnose minnow	208	5.3	0.8 - 2.5	0.31	5.3
Sand shiner	100	2.5	0.8 - 2.3	0.12	2.0
Steelcolor shiner	84	2.1	0.9 - 2.3	0.10	1.7
Western mosquitofish	44	1.1	0.7 - 1.4	0.03	0.5
Smallmouth buffalo	40	1.0	1.1 - 2.8	0.15	2.6
Brook silverside	16	0.4	2.1 - 2.7	0.05	0.9
River carpsucker	10	0.3	1.8 - 2.8	0.05	0.9
Silverjaw minnow	10	0.3	1.0 - 2.1	0.02	0.3
Quillback	9	0.2	1.4 - 2.5	0.04	0.7
Gizzard shad	4	0.1	3.9 - 4.9	0.12	2.0
Striped shiner	4	0.1	2.0 - 2.3	0.02	0.3
Spotted bass	1	*	4.0	0.03	0.5
Golden redhorse	1	*	3.2	0.02	0.3
Central stoneroller	1	*	2.7	0.01	0.2
Rainbow darter	1	*	1.7	0.01	0.2
Total - Species 17	3,927	100.0		5.86	100.0

^{*}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Gizzard shad	53	29.6	4.4 - 13.0	12.69	4.9
River carpsucker	21	11.7	12.9 - 16.0	33.77	13.1
Smallmouth buffalo	12	6.7	18.9 - 24.0	66.46	25.7
Shortnose gar	10	5.6	20.5 - 25.0	14.38	5.6
Spotfin shiner	10	5.6	1.4 - 2.7	0.04	*
Common carp	9	5.0	19.7 - 27.5	56.17	21.7
Channel catfish	9	5.0	11.5 - 20.1	10.93	4.2
Spotted bass	9	5.0	3.4 - 12.3	3.09	1.2
Freshwater drum	6	3.4	12.3 - 15.5	7.09	2.7
Sand shiner	5	2.8	2.1 - 2.4	0.02	*
Steelcolor shiner	4	2.2	1.4 - 3.2	0.03	*
Bigmouth buffalo	3	1.7	22.5	19.06	7.4
Black buffalo	3	1.7	13.7 - 23.5	10.63	4.1
Sauger	3	1.7	14.0 - 15.4	2.90	1.1
White bass	3	1.7	9.2 - 12.9	1.71	0.7
Suckermouth minnow	3	1.7	3.3 - 3.5	0.05	*
Shorthead redhorse	2	1.1	4.6 - 18.8	2.73	1.1
Chestnut lamprey	2	1.1	8.9 - 9.8	0.13	0.1
Longear sunfish	2	1.1	4.0 - 4.3	0.12	*
Mississippi silvery minnow	2	1.1	3.7 - 3.8	0.04	*
Bullhead minnow	2	1.1	2.4 - 2.5	0.01	*
Striped bass	1	0.6	29.5	10.75	4.2
Blue sucker	1	0.6	21.3	2.92	1.1
Quillback	1	0.6	16.2	1.89	0.7
Flathead catfish	1	0.6	13.4	0.92	0.4
Smallmouth bass	1	0.6	7.0	0.16	0.1
Bluntnose minnow	1	0.6	2.6	0.01	*
Total - Species 27	179	100.0		258.70	100.0

^{*}Less than 0.1%.

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	2,071	63.4	0.7 - 2.8	2.82	53.8
Bluntnose minnow	399	12.2	1.0 - 2.8	0.74	14.1
Sand shiner	283	8.7	0.9 - 2.4	0.52	9.9
Bullhead minnow	212	6.5	0.9 - 2.4	0.25	4.8
Steelcolor shiner	80	2.4	0.8 - 2.1	0.11	2.1
Silverjaw minnow	70	2.1	1.4 - 2.4	0.14	2.7
Brook silverside	37	1.1	1.8 - 3.0	0.09	1.7
Smallmouth buffalo	33	1.0	1.6 - 2.7	0.16	3.1
Western mosquitofish	31	0.9	0.6 - 1.9	0.02	0.4
River carpsucker	17	0.5	1.8 - 2.7	0.12	2.3
Quillback	13	0.4	1.5 - 2.8	0.09	1.7
Bluegill	6	0.2	1.0 - 2.2	0.02	0.4
Golden redhorse	4	0.1	3.2 - 3.6	0.07	1.3
Spotted bass	3	0.1	3.5 - 3.6	0.06	1.1
Rosyface shiner	3	0.1	1.5 - 1.8	*	**
Striped shiner	2	0.1	2.2 - 2.3	0.01	0.2
Central stoneroller	1	**	2.6	0.01	0.2
Gizzard shad	1	**	2.4	0.01	0.2
Bigeye chub	1	**	2.2	*	**
Dusky darter	1	**	1.7	*	**
Redfin shiner	1	**	1.7	*	**
Total - Species 21	3,269	100.0		5.24	100.0

^{*}Less than 0.01 lb.

^{**}Less than 0.1%.

NAME OF STREAM: West Fork White River

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Gizzard shad	28	16.3	5.0 - 10.5	4.67	3.0
Suckermouth minnow	26	15.1	3.2 - 3.6	0.42	0.3
Central stoneroller	15	8.7	2.8 - 3.3	0.16	0.1
Channel catfish	14	8.1	4.8 - 25.6	20.71	13.5
River carpsucker	12	7.0	9.9 - 16.3	15.77	10.3
Spotted bass	11	6.4	4.0 - 12.8	6.14	4.0
Spotfin shiner	9	5.2	1.4 - 3.1	0.05	*
Sand shiner	9	5.2	2.0 - 2.3	0.04	*
Smallmouth buffalo	6	3.5	9.4 - 20.6	19.92	13.0
Northern hogsucker	6	3.5	4.0 - 5.4	0.21	0.1
Sauger	5	2.9	12.4 - 18.5	5.92	3.9
Common carp	4	2.3	20.8 - 26.5	28.69	18.7
Golden redhorse	3	1.7	14.2 - 14.4	3.84	2.5
Shorthead redhorse	3	1.7	11.1 - 17.0	3.34	2.2
Smallmouth bass	3	1.7	8.5 - 12.5	1.91	1.2
Logperch	3	1.7	3.8 - 5.8	0.10	0.1
Longnose gar	2	1.2	37.5 - 38.0	10.50	6.8
Bigmouth buffalo	2	1.2	14.2 - 20.2	6.83	4.4
Quillback	2	1.2	14.8 - 17.8	4.44	2.9
Freshwater drum	2	1.2	13.3 - 14.1	1.40	0.9
Flathead catfish	1	0.6	29.2	11.63	7.6
Silver redhorse	1	0.6	22.6	5.15	3.4
Shortnose gar	1	0.6	23.0	1.55	1.0
Bluegill	1	0.6	4.3	0.05	*
Longear sunfish	1	0.6	3.6	0.03	*
Bullhead minnow	1	0.6	2.4	0.01	*
Steelcolor shiner	1	0.6	3.1	0.01	*
Total - Species 27	172	100.0		153.49	100.0

^{*}Less than 0.1%.

NAME OF STREAM: West Fork White River

COMMON NAME	NUMBER	PERCENTAGE	SIZE RANGE (INCHES)	TOTAL WEIGHT (POUNDS)	PERCENTAGE
Spotfin shiner	1,941	80.0	0.6 - 2.7	1.78	53.3
Steelcolor shiner	167	6.9	1.0 - 2.9	0.15	4.5
Sand shiner	127	5.2	1.0 - 2.4	0.22	6.6
Bullhead minnow	88	3.6	0.9 - 2.9	0.07	2.1
Silverjaw minnow	78	3.2	1.2 - 2.6	0.15	4.5
Bluntnose minnow	23	0.9	1.2 - 2.3	0.03	0.9
White bass	1	*	12.4	0.91	27.2
Mississippi silvery minnow	1	*	4.0	0.03	0.9
Total - Species 8	2,426	100.0		3.34	100.0

^{*}Less than 0.1%.

APPENDIX E

STREAM HABITAT EVALUATION FORMS FOR EACH SITE, WFWR 2004

STREAM: W	hite River				RIVER	MILE: 3.6							
NEAREST TO	EAREST TOWN: Mt. Carmel, IL COUNTY: Gibson/Pike NUADRANGLE: East Mount Carmel TWP: 1S RNG: 11W SEC: 18												
QUADRANGL	.E: <u>E</u> 8	ast Mount C	Carmel	TWP:	1S	RNG:	11W	SEC:	18				
LATITUDE:	Upstream 4	253626 N			LONGITUDE: <u>U</u>	Jpstream 1	16 439929 E						
LATITUDE:	Middle 4253	3418 N			LONGITUDE: N	/liddle 16 4	139521 E						
LATITUDE:	Downstrear	n 4253150	N		LONGITUDE: [Downstrea	m 16 439264	4 E					
U.S.G.S. GUA	GING STATIO	N LOCATION	l:	Pet	ersburg	AVO	G. DISCHARGE	(cfs):	2580				
IS REACH RE	PRESENTATI	VE OF STREA	AM (Y/N)	No	IF NOT, WHY? Samp	led below	Kelly's Ripp	le, riffles	are few				
and far bet	ween in this	area											
DESCRIPTIO	N OF SAMPLE	SITE (Access	s, length, di	rection sampled):	Launched at Mt. Ca	rmel, IL ra	mp, motore	d across					
Wabash Ri	ver and up t	he White 3.	.6 miles										
				COLLECTIO	ON SUMMARY								
DATE: 9/16/2004 GEAR: DC Boat Electrofishing EFFORT: 1 hour													
CREW: Stefanavage, Hansen, Ferguson													
OTHER GEAL	R/EFFORT: Se	eine - 4, 50	ft. hauls			WA	TER STAGE:	lo	DW				
CANOPY (%C	DPEN):	100%		PHOTOS (Y/N):	N	_ SECCHI E	OISK (inches):	1	4"				
AIR TEMP (F	:	82.9		WATER TEMP (F):	80.1		D.O. (ppm):		8				
CONDUCTIV	TY:	656	3	pH:	9		ALKALINITY:	13	6.8				
TDS:													
STREAM ME	ASUREMENTS	AVG. WIE	OTH:	506.4	AVG. DEPTH:	38.4	NAX [DEPTH:_	75.6				
STATION LE	NGTH: (1st date	e)		2796	(2nd date)								
WIDTH (ft)		DEPTH (in)											
522	25.2	24	27.6										
483	49.2	42	24		7		7						
522	36	36	49.2		SUBJECTIV	VE /	AESTHETIC						
480	25.2	28.8	75.6		RATING		RATING						
525	31.2	43.2	57.6		(1-10)		(1-10)						
ADDITIONAL (COMMENTS/PC	OLLUTION IMP	ACTS:										
			_										

STREAM: V	/hite River				RIVE	R MILE: 15	5.9		
NEAREST TO	OWN: <u>Hazelt</u>	on, IN			COU	NTY: Gibs	on/Knox		
QUADRANG	LE:	Patoka	1	TWP:	1N	RNG:	11W	SEC:	25
LATITUDE:	Upstream 4	1260778 N			LONGITUDE:	Upstrean	n 16 448631 E	_	
LATITUDE:	Middle 425	9984 N			LONGITUDE:	Middle 16	6 448972 E		
LATITUDE:	Downstrea	m 4259728	N		LONGITUDE:	Downstre	eam 16 44947	7 E	
U.S.G.S. GU	AGING STATIO	ON LOCATION	N:	Pet	ersburg	A	VG. DISCHARG	E (cfs):	2200
IS REACH RI	EPRESENTAT	IVE OF STRE	AM (Y/N)	No	IF NOT, WHY? Sam	pled alon	g north side o	of island a	lthough
south side	was dried u	p. Site had	better ha	bitat than typic	al in area.				
DESCRIPTIO	N OF SAMPLE	E SITE (Acces	s, length, di	rection sampled):	Launched at DFW	public ac	cess site in H	azelton a	nd
motored ~2	2.9 miles do	wnstream to	o island.	Sampled from	head of island dow	nstream,	, 30 minutes a	long eacl	n bank.
				COLLECTIO	ON SUMMARY				
DATE:		9/20/2004		GEAR:	DC Boat Electro	ofishing	EFFORT:	1 h	our
CREW: Stef	fanavage, H	ansen, Ferg	guson						
OTHER GEA	R/EFFORT: S	eine - 4, 50	ft. hauls			W	/ATER STAGE:	lo)W
CANOPY (%	OPEN):	100%		PHOTOS (Y/N):	N	SECCH	II DISK (inches):	1	2"
AIR TEMP (F	·):	76.3		WATER TEMP (F):	76.6		D.O. (ppm):	1	0
CONDUCTIV	TITY:	67	8	pH:	9		ALKALINITY:	1	71
TDS:									
STREAM ME	ASUREMENT	S AVG. WIE	OTH:	329.4	AVG. DEPTH:	69.6	MAX	DEPTH:	144
STATION LE	NGTH: (1st da	te)		4758	(2nd date)				
WIDTH (ft)		DEPTH (in)							
315	33.6	42	48						
339	24	48	61.2		8		8		
267	55.2	82.8	96		SUBJECT	IVE	AESTHETIC		
384	58.8	60	55.2		RATINO	€	RATING		
342	110.4	129.6	144		(1-10)		(1-10)		
ADDITIONAL	COMMENTS/P	OLLUTION IMF	PACTS:	_					

STREAM: W	hite River				RIVER	MILE: <u>24.3</u>	
NEAREST TO	wn: Decke	er, IN			COUNT	Y: Gibson/Knox	
QUADRANGL	E:	Decker	-	TWP:	1N R	RNG: 10W	SEC: 22
LATITUDE:	Upstream 4	4261840 N			LONGITUDE: U	pstream 16 455064 E	
LATITUDE:	Middle 426	1870 N			LONGITUDE: M	iddle 16 454608 E	
LATITUDE:	Downstrea	m 4261271	N		LONGITUDE: De	ownstream 16 45388	7 E
U.S.G.S. GUA	GING STATIO	ON LOCATION	l:	Pet	ersburg	AVG. DISCHARG	E (cfs): 2250
IS REACH RE	PRESENTAT	IVE OF STRE	AM (Y/N)	Yes	IF NOT, WHY?		
DESCRIPTIO	N OF SAMPLE	E SITE (Acces	s, length, di	rection sampled):	Launched at DFW p	ublic access site in H	azelton and
motored ap	prox. 5.6 m	niles upstrea	m. Static	on started at do	ownstream most hou	se along farm land or	n south bank.
Electrofishe	ed for 30 mi	inutes along	each bar	nk.			
				COLLECTIO	ON SUMMARY		
DATE:		9/21/2004		GEAR:	DC Boat Electrofi	ishing EFFORT:	1 hour
CREW: Stef	anavage, H	ansen, Ferg	juson				
OTHER GEAR	R/EFFORT: S	eine - 4, 50	ft. hauls			WATER STAGE:	low
CANOPY (%C	PEN):	100%		PHOTOS (Y/N):	N	SECCHI DISK (inches):	9"
AIR TEMP (F)	:	72.9		WATER TEMP (F):	78.1	D.O. (ppm):	13
CONDUCTIVI	TY:	672	2	pH:	9	ALKALINITY:	153.9
TDS:							
					AVG. DEPTH:	45.6 MAX	DEPTH: 87.6
STATION LEN	IGTH: (1st da	te)		5058	(2nd date)		
WIDTH (ft)		DEPTH (in)		–			
234	43.2	45.6	12	_			
330	70.8	42	24		8	8	
462	87.6	48	18	_	SUBJECTIV		
396	49.2	48	22.8	_	RATING (1-10)	RATING (1-10)	
306	34.8	57.6	72		(1-10)	(1-10)	
ADDITIONAL (COMMENTS/P	OLLUTION IMF	ACTS:				

STREAM: W	hite River				RIVER	MILE: <u>39.2</u>			
NEAREST TO	wn: Peters	burg, IN			COUNT	Y: Knox/P	ike		
QUADRANGL	E:	Monroe C	ity	TWP:	<u>1N</u> F	RNG:	9W	SEC:	12&13
LATITUDE:	Upstream 4	1264324 N			LONGITUDE: U	pstream 1	6 468424 E		
LATITUDE:	Middle 426	3503 N			LONGITUDE: M	liddle 16 4	68507 E		
LATITUDE:	Downstrea	m 4263020	N		LONGITUDE: D	ownstrean	n 16 468280	0 E	
U.S.G.S. GUA	GING STATIO	ON LOCATION	:	Pet	ersburg	AVG	. DISCHARGE	E (cfs):	2200
IS REACH RE	PRESENTAT	IVE OF STREA	AM (Y/N)	Yes	IF NOT, WHY?				
DESCRIPTIO	N OF SAMPLE	E SITE (Access	s, length, di	rection sampled):	Launched at DFW S	R 61 publi	ic access si	te and i	motored
6.5 miles d	ownstream	until stoppe	d by low v	water. Electrof	ished downstream a	t RM 39.2	, 30 minutes	s along	each
bank.									
				COLLECTIO	N SUMMARY				
DATE:	!	9/22/2004		GEAR:	DC Boat Electrof	ishing	EFFORT:	1	hour
CREW: Stef	anavage, H	ansen, Ferg	uson						
OTHER GEAR	R/EFFORT: S	eine - 4, 50	ft. hauls			WAT	ER STAGE:		low
CANOPY (%C	PEN):	100%		PHOTOS (Y/N):	N	SECCHI DI	ISK (inches):		10"
AIR TEMP (F)	:	82.6		WATER TEMP (F):	79.2		D.O. (ppm):		15
CONDUCTIVI	TY:	686	3	pH:	9	A	LKALINITY:	1	53.9
TDS:									
					AVG. DEPTH:	54	MAX I	DEPTH:	118.8
STATION LEI	NGTH: (1st da	te)	ı	4740	(2nd date)				
\\\\\DTU (#\)		DEDTH (in)							
WIDTH (ft)	64.2	DEPTH (in)	40	$\overline{\ }$					
546	61.2	69.6	48		7				
324	76.8	64.8	12			·	7		
297	118.8	85.2	18		SUBJECTIV RATING	E A	ESTHETIC RATING		
417	81.6	90	18	-	(1-10)		(1-10)		
636	27.6	24	18						
ADDITIONAL (COMMENTS/PO	OLLUTION IMP	ACTS:						

STREAM: W	/hite River				RIVER	MILE: 45.7						
NEAREST TO	EAREST TOWN: Petersburg COUNTY: Pike/Knox UADRANGLE: Monroe City TWP: 1N RNG: 8W SEC: 15											
QUADRANG	_E:	Monroe (City	TWP:	<u>1N</u> R	NG: 8W	SEC: 15					
LATITUDE:	Upstream 4	4262573 N			LONGITUDE: U	ostream 16 474840 E						
LATITUDE:	Middle 426	2589 N			LONGITUDE: M	iddle 16 474560 E						
LATITUDE:	Downstrea	m 4262429	N		LONGITUDE: Do	ownstream 16 47432	7 E					
U.S.G.S. GU	AGING STATIO	ON LOCATION	N:	Pete	ersburg	AVG. DISCHARGE	E (cfs): 2150					
IS REACH RI	EPRESENTAT	IVE OF STRE	AM (Y/N)	Yes	IF NOT, WHY?							
DESCRIPTIO	N OF SAMPLI	E SITE (Acces	s, length, di	rection sampled):	Launched at DFW bo	oat ramp at SR 61 bri	dge.					
Electrofish	ed from brid	lge downstr	eam to Pr	ide's Creek, 30	minutes along each	ı bank.						
				COLLECTIO	N SUMMARY							
DATE:		9/23/2004		GEAR:	DC Boat Electrofi	shing EFFORT:	1 hour					
CREW: Stef	anavage, H	ansen, Ferç	guson									
OTHER GEA	R/EFFORT: S	eine - 4, 50	ft. hauls			WATER STAGE:	low					
CANOPY (%	OPEN):	100%		PHOTOS (Y/N):	N	SECCHI DISK (inches):	13"					
AIR TEMP (F):	67.5		WATER TEMP (F):	80.6	D.O. (ppm):	8					
CONDUCTIV	ITY:	70	2	pH:	9	ALKALINITY:	153.9					
TDS:												
STREAM ME	ASUREMENT	S AVG. WII	DTH:	481.8	AVG. DEPTH:	64.8 MAX	DEPTH: 140.4					
STATION LE	NGTH: (1st da	te)		1794	(2nd date)							
WIDTH (ft)		DEPTH (in)		7								
486	140.4	44.4	24	_								
399	91.2	72	66	_	7	7						
450	97.2	92.4	67.2	_	SUBJECTIVI							
522	57.6	54	56.4	_	RATING (1-10)	RATING (1-10)						
552	43.2	27.6	38.4	4	, ,	,						
ADDITIONAL	COMMENTS/P	OLLUTION IMF	PACTS:									

STREAM: W	est Fork WI	hite River			RI\	ER MILE:	57.5		
NEAREST TO	wn: Washii	ngton			cc	OUNTY: Da	aviess/Knox		
QUADRANGL	E:	Sandy Ho	ook	TWP:	3N, 4N	RNG:	7W, 8W	SEC:	7, 18
LATITUDE:	Upstream 3	88.61747° N	I		LONGITUD	E: Upstre	am -87.23979° \	N	
LATITUDE:	Downstrear	n 38.60852	.° N		LONGITUD	E: Downs	tream -87.24619	9° W	
U.S.G.S. GUA	GING STATIC	N LOCATION	J:	Ne	ewberry		AVG. DISCHARGI	E (cfs):	3,400
IS REACH RE	PRESENTATI	VE OF STRE	AM (Y/N)	Yes	IF NOT, WHY?				
DESCRIPTION	N OF SAMPLE	SITE (Acces	s, length, d	irection sampled):	Started at a set of	of house	foundation poles	on east	bank.
Launched a	at Washingto	on ramp							
				COLLECTIO	ON SUMMARY				
DATE:	ę	9/21/2004		GEAR:	DC EF	Boat	EFFORT:	1	HR
CREW: Dan	Carnahan,	Jason Doll,	Brant Fis	sher					
OTHER GEAR	R/EFFORT: 3	Seine hauls	5				WATER STAGE:	2.4	45 ft
CANOPY (%C)PEN):	100		PHOTOS (Y/N):	No	SEC	CHI DISK (inches):	1	0.0
AIR TEMP (F)	:	59.9		WATER TEMP (F):	68.2	2	D.O. (ppm):	1	1.6
CONDUCTIVI	TY:	827	μs	pH:	7.72		ALKALINITY:	205.	2 ppm
TDS:									
STREAM MEA	ASUREMENTS	S AVG. WI	OTH:	318.6 ft	AVG. DEPTH:	36.4	4 in MAX	DEPTH:_	81.6 in
STATION LEN	NGTH: (1st dat	e)	5	5,050 ft	(2nd da	ite)			
144DTH (6)		DEDTIL (1)							
WIDTH (ft)	40.0	DEPTH (in)	00.0	7					
	12.0 25.2	24.0	36.0		Г		4		
		51.6	50.4	-			· <u></u>		
300.0	12.0	36.0	81.6	\dashv	SUBJE:		AESTHETIC RATING		
387.0	12.0	27.6	81.6	\dashv	(1-1		(1-10)		
297.0	33.6	39.6	22.8	_					
ADDITIONAL (COMMENTS/PC	OLLUTION IMF	PACTS:						

STREAM: W	est Fork W	hite River			R	IVER MILE	E: <u>61.52</u>	
NEAREST TO	วพท: <u>Washi</u> เ	ngton			C	OUNTY: <u>I</u>	Daviess, Knox	
QUADRANGL	.E:	Washingt	on	TWP:	3N	RNG	8W, 7W	SEC: 62, 295
LATITUDE:	Upstream 3	88.65435° N			LONGITUI	DE: Upstr	ream -87.24036°\	N
LATITUDE:	Downstrear	n 38.64448	° N		LONGITUI	DE: <u>Dowr</u>	nstream -87.23838	3° W
U.S.G.S. GUA	AGING STATIC	N LOCATION	l:	Ne	ewberry		AVG. DISCHARG	E (cfs): 3,400
IS REACH RE	EPRESENTATI	VE OF STREA	AM (Y/N)	Yes	IF NOT, WHY?			
DESCRIPTIO	N OF SAMPLE	SITE (Access	s, length, di	irection sampled):	Launched at W	ashingto	n ramp. Station e	ended just
downstrear	m of SR 50 E	Bridge.						
				COLLECTIO	ON SUMMARY			
DATE:	(9/21/2004		GEAR:	DC EF	Boat	EFFORT:	1 HR
CREW: Dan	Carnahan,	Jason Doll,	Brant Fis	sher				
OTHER GEAL	R/EFFORT: 3	Seine hauls	1				WATER STAGE:	2.45 ft
CANOPY (%C	OPEN):	100		PHOTOS (Y/N):	No	SE	CCHI DISK (inches):	10.0
AIR TEMP (F):	59.9		WATER TEMP (F):	68.	.2	D.O. (ppm):	11.6
CONDUCTIV	ITY:	803	μs	pH:	7.7		ALKALINITY:	205.2 ppm
TDS:								
STREAM ME	ASUREMENTS	AVG. WIE	OTH:	301.0 ft	AVG. DEPTH:	49	9.2 inMAX	DEPTH: 111.6 in
STATION LE	NGTH: (1st dat	e)	3	3,804 ft	(2nd c	date)		
WIDTH (ft)	22.2	DEPTH (in)	00.0					
228.0		76.8	82.8			4	4	
	60.0	48.0	33.6	_				
321.0	33.6	45.6	30.0	_		ECTIVE TING	AESTHETIC RATING	
336.0	25.2	42.0	12.0	_		-10)	(1-10)	
303.0	111.6	123.6	6.0	\dashv				
312.0	34.8	31.2	28.8					
ADDITIONAL (COMMENTS/PC	DLLUTION IMP	ACTS:					

STREAM: W	est Fork WI	nite River				RIVER MILI	E: 73.06			
NEAREST TO	WN: Bickne	II				COUNTY:	Daviess, Knox			
QUADRANGL	E:	Wheatlar	nd	TWP:	3N, 4N	RNG	9W	SEC:	1, 36	
LATITUDE:	Upstream 3	8.75030° N			LONGITU	JDE: Upst	ream -87.24031° V	V		
LATITUDE:	Downstrear	n 38.74837°	, N		LONGITU	JDE: Dow	nstream -87.24663	3° W		
U.S.G.S. GUA	GING STATIO	N LOCATION:		Ne	ewberry		AVG. DISCHARGE	E (cfs):	3,400	
IS REACH RE	PRESENTATI	VE OF STREA	M (Y/N)	Yes	IF NOT, WHY?					
DESCRIPTION	ON OF SAMPLI	E SITE (Access	s, length, direc	ction sampled):	Launched at C	Carnahan	ramp, went downs	tream.		
				COLLECTIO	ON SUMMARY	•				
DATE:	(9/22/2004		GEAR:	DC E	F Boat	EFFORT:		I HR	
CREW: Dan	Carnahan,	Jason Doll,	Brant Fish	er						
OTHER GEAF	OTHER GEAR/EFFORT: 3 Seine hauls WATER STAGE: 2.45 ft									
CANOPY (%C	PEN):	100	P	HOTOS (Y/N):	No	SI	ECCHI DISK (inches):		11.0	
AIR TEMP (F)	:	64.9	w	/ATER TEMP (F):	69	9.4	D.O. (ppm):		11.4	
CONDUCTIVI	TY:	874	JS	pH:	7.7		ALKALINITY:	239).4 ppm	
TDS:										
STREAM MEA	SUREMENTS	AVG. WID	TH:2	217.5 ft	AVG. DEPTH:	5	5.3 in MAX	DEPTH:	264 in	
STATION LEN	IGTH: (1st date	e)	3,5	11 ft	(2nd	date)				
WIDTH (ft)		DEPTH (in)		1						
189.0		135.6			Γ					
297.0	50.4	90.0	12.0			8	8			
168.0	99.6	79.2	12.0	-		JECTIVE ATING	AESTHETIC RATING			
213.0	146.6	110.4	12.0			1-10)	(1-10)			
246.0	6.0	12.0	14.4	-						
192.0	31.2	12.0	6.0	J						
ADDITIONAL C	OMMENTS/PO	LLUTION IMPA	CTS:							

STREAM: W	est Fork W	hite River			RIVER	MILE: <u>84.5</u>	
NEAREST TO	own: Edwar	dsport			COUNT	Y: Daviess, Knox	
QUADRANGL	.E:	Plainville	e	TWP:	4N F	RNG: 8W, 7W	SEC: 1, 6
LATITUDE:	Upstream 3	38.81374° N			LONGITUDE: U	pstream -87.23320°	W
LATITUDE:	Downstrea	m 38.81001	° N		LONGITUDE: D	ownstream -87.2447	7° W
U.S.G.S. GUA	AGING STATIO	ON LOCATION	l:	Ne	ewberry	AVG. DISCHARG	E (cfs): 3,400
IS REACH RE	EPRESENTAT	IVE OF STREA	AM (Y/N)	Yes	IF NOT, WHY?		
DESCRIPTIO	N OF SAMPLE	SITE (Access	s, length, dir	ection sampled):	Station ended under	high power lines.	
Launched a	at Carnahan	boat ramp.					
				COLLECTIO	ON SUMMARY		
DATE:	!	9/20/2004		GEAR:	DC EF Boa	t EFFORT:	1 HR
CREW: Dan	Carnahan,	Jason Doll,	Brant Fis	her			
OTHER GEAL	R/EFFORT: 3	Seine hauls	1			WATER STAGE:	2.45 ft
CANOPY (%0	OPEN):	100		PHOTOS (Y/N):	No	SECCHI DISK (inches):	11.0
AIR TEMP (F):	64.9		WATER TEMP (F):	69.4	D.O. (ppm):	11.4
CONDUCTIV	ITY:	874	μs	pH:	7.7	ALKALINITY:	239.4 ppm
TDS:							
STREAM ME	ASUREMENTS	S AVG. WIE	OTH:	289.0 ft	AVG. DEPTH:	45.3 in MAX	DEPTH: 100.8 in
STATION LE	NGTH: (1st da	te)	4,	803 ft	(2nd date)		
WIDTH (ft)		DEPTH (in)					
	87.6		6.0	\exists			
		67.2		1	5	5	
306.0	36.0	40.8	18.0	1	SUBJECTIV		
213.0	6.0	49.2	88.8	1	RATING		
285.0	18.0	36.0	70.8		(1-10)	(1-10)	
339.0	64.8	38.4	6.0	1			
	<u> </u>	OLLUTION IMP		- 1			
, DELITORAL (O O IVIIVILIA I O/F	OLLO HON HVIF					

STREAM: W	est Fork W	hite River			RIVER	MILE: 101.4					
NEAREST TO	wn: Elnora				COUNT						
QUADRANGL	.E: Lyon	ns, Epsom,	Plainville	TWP:	5N F	RNG: 6W	SEC	18			
LATITUDE:	Upstream 3	38.87656° N			LONGITUDE: U	pstream -87.1223	32° W				
LATITUDE:	Downstrea	m 38.87019	° N		LONGITUDE: D	ownstream -87.12	2805° W				
U.S.G.S. GUA	AGING STATIO	ON LOCATION	l:	Ne	ewberry	AVG. DISCHA	ARGE (cfs):	3,400			
IS REACH RE	EPRESENTAT	IVE OF STREA	AM (Y/N)	Y	IF NOT, WHY?						
DESCRIPTIO	N OF SAMPLE	SITE (Access	s, length, dir	ection sampled):	Station started just of	downstream of SR	58 Bridge	and			
went down	stream. Lau	unched at E	Inora ram	р.							
				COLLECTIO	ON SUMMARY						
DATE:		9/20/2004		GEAR:	DC EF Boa	t EFFO	RT:	1 HR			
CREW: Dan	Carnahan,	Jason Doll,	Brant Fis	her							
OTHER GEAL	OTHER GEAR/EFFORT: 3 Seine hauls WATER STAGE: 2.45 ft										
CANOPY (%0	OPEN):	100		PHOTOS (Y/N):	No	SECCHI DISK (inch	es):	12.0			
AIR TEMP (F):	62.8		WATER TEMP (F):	67.8	D.O. (pp	m):	9.1			
CONDUCTIV	TY:	870	μs	pH:	7.9	ALKALINIT	Y: <u>25</u>	6.5 ppm			
TDS:											
STREAM ME	ASUREMENTS	S AVG. WIE	OTH:	223.5 ft	AVG. DEPTH:	33.7 in	MAX DEPTH:	102 in			
STATION LE	NGTH: (1st dat	te)	5,	870 ft	(2nd date)						
WIDTH (ft)		DEPTH (in)									
267.0	36.0		24.0	7							
243.0	24.0	24.0	24.0		6	6					
153.0	48.0	36.0	12.0		SUBJECTIV	ب	TC.				
156.0	60.0	27.0	12.0		RATING						
249.0	102.0	42.0	12.0	1	(1-10)	(1-10)					
273.0	55.5	36.0	12.0	1							
	<u>, </u>	OLLUTION IMP		= 1							
, DELITORAL (JOININE IN 10/F	OLLO HON HVIF									

STREAM: W	est Fork W	hite River			RI	VER MILE:	105.7		
NEAREST TO	wn: Elnora	l			C(DUNTY: Da	aviess/Knox/Gre	ene	
QUADRANGL	.E:	Scotlan	d	TWP:	6N, 5N	RNG:	6W	SEC:	5, 32
LATITUDE:	N 38.90899	9			LONGITUD	E: W 87.0)9615		
LATITUDE:	N 38.89955	5			LONGITUD	DE: W 87.1	10472		
U.S.G.S. GUA	AGING STATIO	ON LOCATION	l:	Ne	wberry		AVG. DISCHARGE	E (cfs):	908
IS REACH RE	EPRESENTAT	IVE OF STRE	AM (Y/N)	Υ	IF NOT, WHY?				
DESCRIPTIO	N OF SAMPLE	E SITE (Acces	s, length, dire	ction sampled):	Launched at DN	IR ramp N	NW of Elnora. Sa	ampled ju	ust
				p, top of san					
	g-			p, 10 p 11 10					
				COLLECTIO	N SUMMARY				
DATE:	!	9/20/2004		GEAR:	DC B	oat	EFFORT:	1	hr.
CREW: Kitta	ıka, King, La	ang							
OTHER GEAL	R/EFFORT: S	eine (4)					WATER STAGE:	No	rmal
CANOPY (%0	OPEN):	90	P	HOTOS (Y/N):	Υ	SEC	CHI DISK (inches):	1	16"
AIR TEMP (F):	75	w	/ATER TEMP (F):	72		D.O. (ppm):	1	4.5
CONDUCTIV	TY:			pH:	9.5		ALKALINITY:	11	19.7
TDS:									
STREAM ME	ASUREMENTS	S AVG. WIE	OTH:	288'	AVG. DEPTH:	58	B"MAX	DEPTH:_	156"
STATION LE	NGTH: (1st da	te)	427	71 ft.	(2nd da	ate)			
WIDTH (ft)		DEPTH (in)							
216	145	110	63	1					
324	40	25	17	1	[5	3		
342	16	24	47	1	_	CTIVE	AESTHETIC		
261	24	27	45	1	RAT	ING	RATING		
306	348	105	156		(1-	10)	(1-10)		
ADDITIONAL (COMMENTS/P	OLLUTION IMF	PACTS: Lau	nched NW o	of Elnora. 1550N	to 400E,	follow to public a	access s	ite
				d boulders.					
								_	

STREAM: W	est Fork Wh	ite River		RIVER N	RIVER MILE: 115.5					
NEAREST TO	OWN: Plumm	er				COUNT	Y: Greer	ne		
QUADRANGL	.E:	Scotland	d	TWP:	16	N R	NG:	5W	SEC:	17
LATITUDE:	N38.95869				L	ONGITUDE: W	86.9932	6		
LATITUDE:	N38.95195				L	ONGITUDE: W	86.9958	3		
U.S.G.S. GUA	AGING STATION	N LOCATION:		Ne	ewberry	G. DISCHARGE	E (cfs):	908		
IS REACH RE	EPRESENTATIV	E OF STREA	M (Y/N) _	Y	IF NOT, V	VHY?				
DESCRIPTION	ON OF SAMPLE	SITE (Access	s, length, d	irection sampled):	Started	at Nedich Dit	tch and e	ended above	Doans	Creek.
				COLLECTIO	ON SUM	MARY				
DATE:	9	/23/2004		GEAR:		DC Boat		EFFORT:	1	hr.
CREW: Kitta	aka, King, La	ng								
OTHER GEAR/EFFORT: Seine (4)							WA	TER STAGE:	No	rmal
CANOPY (%OPEN):				PHOTOS (Y/N):		Υ	SECCHI	DISK (inches):		18"
AIR TEMP (F)):	82·		WATER TEMP (F):	D.C			D.O. (ppm):		17
CONDUCTIV	ITY:	1.0°	1	pH:	pH: 9.2			ALKALINITY:	1:	36.8
TDS:					0.5					
STREAM ME	ASUREMENTS	AVG. WID	TH:	320.4'	AVG. DE	PTH:	42.5"	MAX I	DEPTH:_	62"
STATION LE	NGTH: (1st date))		3084'		(2nd date)				
WIDTH (ft)		DEPTH (in)								
324	141	54	38							
315	150	41	45			5		4		
318	96	44	62			SUBJECTIVE	≣	AESTHETIC		
312	165	36	33			RATING		RATING		
333	75	40	36			(1-10)		(1-10)		
ADDITIONAL (COMMENTS/POL	LUTION IMPA	стѕ: L	aunched at pri	vate prop	perty under c	overed b	oridge over P	<u>lum</u> mer	Creek.
Permission	n from Jerry L	_ester, 812-	- 384-830	<u></u> 14.						
_						_	_			_

STREAM: W	STREAM: West Fork White River RIVER MILE: 125.0								
NEAREST TO	WN: Bloom	field				COUNTY: Gre	eene		
QUADRANGL	.E:	Bloomfie	eld	TWP:	7N	RNG:	5W	SEC:	21, 28
LATITUDE:	N39.0193				LONGITU	JDE: <u>W86.97</u>	225		
LATITUDE:	N39.03051				LONGITU	JDE: <u>W86.96</u>	651		
U.S.G.S. GUA	AGING STATIO	ON LOCATION	l:				AVG. DISCHARGE	E (cfs):	908
IS REACH RE	EPRESENTAT	IVE OF STRE	AM (Y/N)	Υ	IF NOT, WHY?				
DESCRIPTIO	N OF SAMPLI	E SITE (Acces	s, length, di	rection sampled):	Started above	old bridge p	osts, then dowr	nstream	past the
island.									
				00115071	201 21 10 10 10 10 10 10 10 10 10 10 10 10 10	,			
					ON SUMMARY				
	ATE: 9/21/2004 CREW: Kittaka, King, Lang			GEAR:	DC	Boat	EFFORT:	1	hr.
							WATER STAGE:		
CANOPY (%OPEN): 90			PHOTOS (Y/N):		SECC	HI DISK (inches):		12"	
AIR TEMP (F):	71		WATER TEMP (F):		70	D.O. (ppm):		10
CONDUCTIV	TY:	1.0	1	pH:	9.5		ALKALINITY:		
TDS:					51				
STREAM ME	ASUREMENT	S AVG. WIE	OTH:	275.4 ft.	AVG. DEPTH:	55.3	MAX	DEPTH:_	120"
STATION LE	NGTH: (1st da	te)	4,	351 ft.	(2nd	date)			
WIDTH (ft)		DEPTH (in)							
195	111	92	94						
255	108	31	71		ſ				
321	300	79	24		SUB	JECTIVE	AESTHETIC		
333	312	30	0			ATING	RATING		
273	87	26	76		(1-10)	(1-10)		
ADDITIONAL (COMMENTS/P	OLLUTION IMP	PACTS: La	≕ aunched on N\	N side of river	off HWY 23	1 from River Ro	ad. Priv	ate ramn
		ssion from I				J			ato ramp
at Lattao O			- 3. 5 1 1111,	231 0000.					

STREAM: W	REAM: West Fork White River RIVER MILE: 134.7									
NEAREST TO	WN: Worth	ington			C	OUNTY: Gre	ene			
QUADRANGL	.E:	Bloomfie	eld	TWP:	8N	RNG:	5W	SEC:	21	
LATITUDE:	N39.11676	5			LONGITUE	E: W86.962	244			
LATITUDE:	N39.11034	ļ			LONGITUE	E: W86.963	353			
U.S.G.S. GUA	AGING STATIO	ON LOCATION	N:				AVG. DISCHARG	E (cfs):	908	
IS REACH RE	EPRESENTAT	IVE OF STRE	AM (Y/N)	Υ	IF NOT, WHY?					
DESCRIPTIO	N OF SAMPLI	E SITE (Acces	s, length, dir	ection sampled):	Began sampling	j just above	confluence w	th Eel R	iver,	
downstrear	n to HWY 1	57 bridge.								
				COLLECTIO	ON CHMMARY					
					ON SUMMARY					
				GEAR:	DC B	oat	EFFORT:	1	hr.	
CREW: Kitta										
OTHER GEAR/EFFORT: Seine (4)										
CANOPY (%OPEN):										
					72					
CONDUCTIV	TY:			pH:	10		ALKALINITY:			
TDS:										
STREAM ME	ASUREMENT	S AVG. WII	OTH:	232'	AVG. DEPTH:	63.5	"MAX	DEPTH:_	128"	
STATION LE	NGTH: (1st da	te)	2	2556'	(2nd d	ate)				
WIDTH (ft)		DEPTH (in)								
219	231	85	45	7						
324	147	29	34	1	Γ					
222	87	63	117		SUBJE	ECTIVE	AESTHETIC			
169	63	70	114			ING	RATING			
179	135	35	45		(1-	10)	(1-10)			
159	384	133	44							
ADDITIONAL (COMMENTS/P	OLLUTION IMF	PACTS: La	— iunched at DN	NR ramp on NE s	side of HW	7 157. In good	l conditio	on.	
Very shallo							<u> </u>			

STREAM: W	REAM: West Fork White River RIVER MILE: 145.1									
NEAREST TO	งพท: <u>Farme</u>	rs				COUNTY	: Owen			
QUADRANGL	.E:	Freedor	n	TWP:	9N	RI	NG:	4W	SEC:	33
LATITUDE:	N39.18048				LONG	GITUDE: W	86.85218	3		
LATITUDE:	N39.17346				LONG	GITUDE: W	86.88833	3		
U.S.G.S. GUA	AGING STATIO	ON LOCATION	l:				AV0	G. DISCHARGE	(cfs):	837
IS REACH RE	EPRESENTAT	IVE OF STREA	AM (Y/N)	Y	IF NOT, WH	Y?				
DESCRIPTIO	N OF SAMPLE	SITE (Access	s, length, dir	rection sampled):	Launched a	at HWY 35	8 ramp.	Started san	npling at	the
mouth of R	mouth of Raccoon Creek.									
				COLLECTIO						
DATE:	!	9/22/2004		GEAR:	[DC Boat		EFFORT:	1	hr.
CREW: Kittaka, King, Lang										
OTHER GEAR/EFFORT: Seine (4)							WA	TER STAGE:	No	rmal
CANOPY (%C	DPEN):			PHOTOS (Y/N):			SECCHI D	ISK (inches):	1	3"
AIR TEMP (F)):	80		WATER TEMP (F):		72		D.O. (ppm):		11
CONDUCTIVI	TY:	1.1	6	pH:	!	9.0		ALKALINITY:	1	71
TDS:					0.56					
STREAM MEA	ASUREMENTS	S AVG. WIE	OTH:	220.2'	AVG. DEPTI	H:	45.1"	MAX I	DEPTH:_	145"+
STATION LEN	NGTH: (1st dat	te)	3	3010'		(2nd date)				
WIDTH (ft)		DEPTH (in)								
222	48	28	29	$\overline{\ \ }$						
210	57	38				7		7		
198	60	55	57	1	ç	SUBJECTIVE	: <i>1</i>	AESTHETIC		
222	165	34	17	1		RATING	- <i>'</i>	RATING		
177	81	145+	80	1		(1-10)		(1-10)		
177	01	1401	00	1						
ADDITIONAL (COMMENTS/DA		ACTS: I a	ച nunched at Fa	rmers Ferry	Old Fern	v croeein	a with concr	ete elah	to
				231. Needs cle					cie siau	10
iauricii Al	. Cha di cou	iny isa aad t	711 1 1 V V I Z	-0 1. INCCUS CIC	Janea up-III	oc site iti	ואוזומו	π.		

STREAM: W	REAM: West Fork White River RIVER MILE: 156.36									
NEAREST TO	WN: Potters	sville, Indiar	na		COL	JNTY: Owen				
QUADRANGL	E: Freedom	, IN 1956. Pho	otorevised 19	980 TWP:	9N	RNG:	4W	SEC:	1, 12	
LATITUDE:*	39.24711°	Ma	ap Datum	= NAD83	LONGITUDE	: 86.79943°	Map Da	atum = N	NAD83	
LATITUDE:**	39.23765°	Ma	ap Datum	= NAD83	LONGITUDE	: 86.80122°	Map Da	atum = N	NAD83	
U.S.G.S. GAU	GING STATIO	N LOCATION:		RM 162.21 at	Spencer, Indiana	AVG	. DISCHARGE	E (cfs):		
IS REACH RE	PRESENTATI	VE OF STREA	.M (Y/N)	Υ	IF NOT, WHY?					
DESCRIPTION	ON OF SAMPLI	E SITE (Acces	s, length, dire	ection sampled):	Launched from pr	rivate boat rai	mp on left l	oank in s	ection	
12. Sample	e site starte	d at mouth o	of intermit	tent stream o	n left bank in secti	on 1 approxir	nately 3,40	0 feet		
upstream of boat ramp. Left and right banks were sampled in downstream direction.										
	DATE: 9/21/2004 GEAR: 16-foot DC shocker boat EFFORT: 3,600 seconds									
DATE:	(9/21/2004		GEAR:	16-foot DC sho	cker boat	EFFORT:	3,600	seconds	
CREW: L. Lehman, C. Kowalik, D. Ort										
OTHER GEAR/EFFORT: 4 hauls with 6' X 50' seine (1/8" Delta mesh) WATER STAGE:										
CANOPY (%C	CANOPY (%OPEN):95 PHOTOS (Y/N):Y SECCHI DISK (inches):21									
AIR TEMP (F)	<u> </u>	75		WATER TEMP (F):	68		D.O. (ppm):		12	
CONDUCTIVI	TY:	700 micror	mhos/cm	pH:	8.3	AI	LKALINITY:	274 to	291 ppm	
TDS:			(conduc	ctivity) (0.84 c	orrelation factor) =	588 ppm				
STREAM MEA	SUREMENTS	AVG. WIE	DTH:	190 feet	AVG. DEPTH:	51 inches	MAX	DEPTH:	10 feet	
STATION LEN	IGTH: (1st date	e)	approxima	itely 3,850 fee	t (2nd date	e)	NA	١		
WIDTH (ft)		DEPTH (in)		\exists						
189	72	50		_		1				
201	53	73	48	-]				
144	22	41	79	-	SUBJEC		ESTHETIC			
210	79	48	24	4	RATIN (1-10		RATING (1-10)			
207	37	49	72	_	(1.1.	,	(* **)			
ADDITIONAL C	OMMENTS/PO	LLUTION IMPA	.стs: <u>*Т</u>	op of station (approximately RN	1 156.36). **E	Bottom of s	tation.		
Latitude and longitude near center of station at river mile 156.00 = 39.24230° and 86.79882°.										
Three fish lice were collected and preserved										

STREAM: W	EAM: West Fork White River RIVER MILE: 164.53										
NEAREST TO	wn: Spend	er, Indiana.	Romona	, Indiana	COU	NTY: Owen					
QUADRANGL	.E:	Gosport, IN	1998	TWP:	10N	RNG:	3W	SEC: 15,21, 22			
LATITUDE:*	39.29729°	Ma	ap Datum	= NAD83	LONGITUDE:	86.73784°	Map D	atum = NAD83			
LATITUDE:**	39.29292°	Ma	ap Datum	= NAD83	LONGITUDE:	86.74622°	Map D	atum = NAD83			
U.S.G.S. GAL	JGING STATIO	ON LOCATION	:	RM 162.21 at	Spencer, Indiana	AVG	. DISCHARG	E (cfs):			
IS REACH RE	EPRESENTAT	IVE OF STREA	AM (Y/N)	Υ	F NOT, WHY?			_			
DESCRIPTIO	N OF SAMPLE	E SITE (Access	s, length, dir	rection sampled): L	aunched from IDI	NR ramp at	Spencer. S	Sample site			
extended approximately 3,060 feet downstream (determined with GPS unit) from mouth of McCormicks Creek at											
River Mile	164.53. Lef	t and right b	anks wer	e sampled in do	ownstream direction	on.					
	COLLECTION SUMMARY										
DATE:		9/20/2004		GEAR:	16-foot DC shoo	ker boat	_ EFFORT:	3,600 seconds			
CREW: L. Lehman, C. Kowalik, D. Ort											
OTHER GEAR/EFFORT: 4 hauls with 6' X 50' seine (1/8" Delta mesh) WATER STAGE:											
CANOPY (%C	CANOPY (%OPEN):90 PHOTOS (Y/N):Y SECCHI DISK (inches):20										
AIR TEMP (F)	:	70		WATER TEMP (F):	68		D.O. (ppm):	8 to 9			
CONDUCTIVI	TY:	1,000 micro	mhos/cm	pH:	8.3	A	LKALINITY:	222 to 239 ppm			
TDS:			(condu	ctivity) (0.84 co	rrelation factor) =	840 ppm					
STREAM ME	ASUREMENT	S AVG. WIE)TH:	187 feet	AVG. DEPTH:	61 inches	MAX	DEPTH: >11 feet			
STATION LE	NGTH: (1st da	te)	pproxima	ately 3,060 feet	(2nd date)	NA	1			
WIDTH (ft)		DEPTH (in)		7							
144	50	42	34	-							
237	58	53	61	_							
225	58	66	72	4	SUBJECT		ESTHETIC				
195	42	30	17	_	RATIN((1-10)		RATING (1-10)				
132	>132	120	74	_	(1.10)		(1.13)				
ADDITIONAL (COMMENTS/P	OLLUTION IMP	ACTS: *T	op of station at mo	uth of McCormicks Cr	eek (RM 164.5	3). **Bottom	of station. Latitude			
and longitude	near center of	station at RM	164.24 = 39	9.29474° and 86.74	154°. Remnants of o	ld railroad bridg	ge are presen	t in river 460			
feet upstream	feet upstream of mouth of McCormicks Creek. This shoreline angling site is accessible from Trail 7 in McCormick's Creek State Park.										

STREAM: W	TREAM: West Fork White River RIVER MILE: 194.09										
NEAREST TO	WN: Martin	sville, India	na		COL	JNTY: Morgan					
QUADRANGL	E: Ma	artinsville, IN	N 1998	TWP:	12N	RNG:	1E	SEC:	20, 29		
LATITUDE:*	39.46133°	Ma	ap Datum =	NAD83	LONGITUDE	86.44152°	Map Da	atum = I	NAD83		
LATITUDE:**	39.45371°	Ma	ap Datum =	NAD83	LONGITUDE	86.43811°	Map Da	atum = I	NAD83		
U.S.G.S. GAU	GING STATIO	N LOCATION:	RM	l 198.92 near	Centerton, Indiar	na AVG.	DISCHARGE	E (cfs):			
IS REACH RE	PRESENTATI	VE OF STREA	.M (Y/N)	Υ	IF NOT, WHY?						
DESCRIPTION	ON OF SAMPL	E SITE (Acces	s, length, direc	tion sampled):	Launched from pr	ivate boat rar	np at bait s	shop on	right		
bank. Station started at box culvert on right bank approximately 1,215 feet above boat ramp. Station length of											
3,060 feet o	determined	with GPS u	nit. Left an	d right banks	were sampled in	downstream	direction.				
COLLECTION SUMMARY											
DATE:	,	9/28/2004		GEAR:	16-foot DC sho	cker boat	EFFORT:	3,600	seconds		
CREW: L. Lehman, C. Kowalik, A. Richards											
OTHER GEAR/EFFORT: 4 hauls with 6' X 50' seine (1/8" Delta mesh) WATER STAGE:											
CANOPY (%C	CANOPY (%OPEN): 90 PHOTOS (Y/N): Y SECCHI DISK (inches): 21										
AIR TEMP (F)	:	64	w	ATER TEMP (F):	70		D.O. (ppm):	1:	3-14		
CONDUCTIVI	TY:	1,000 micro	mhos/cm	pH:	8.3	AL	KALINITY:	239 to	256 ppm		
TDS:			(conducti	vity) (0.84 cc	rrelation factor) =	840 ppm					
STREAM MEA	SUREMENTS	AVG. WIE	OTH: 2	34 feet	AVG. DEPTH:	41 inches	MAX	DEPTH:	8 feet		
STATION LEN	IGTH: (1st date	e)	approximate	ely 3,060 feet	(2nd date	e)	NA	1			
WIDTH (ft)		DEPTH (in)		1							
246	52	38	28			1					
231	53	31	28								
258	30	26	26		SUBJEC		ESTHETIC				
165	35	72	76		RATIN (1-10		RATING (1-10)				
270	67	25	28		(,	(1.10)				
				<u>]</u>							
ADDITIONAL COMMENTS/POLLUTION IMPACTS: *Top of station (approximately RM 194.09) at box culvert under SR 67.											
**Bottom of	station. La	atitude and I	ongitude ne	ear center of	station at RM 193	3.80 = 39.457	36° and 86	.44075°			
Latitude and longitude at boat ramp = 39.45796° and 86.44147°.											

APPENDIX F

QHEI FORMS FOR EACH SITE, WFWR 2004

STREAM: White River	RIVER MILE	3.6	DATE:	9/16/2004	QHEI SCORE 46	6
1) SUBSTRATE: (Check ONLY Two Substrate TYPE POOL RIFFLE BLDER/SLAB(10) BOULDER(9) COBBLE(8) X HARDPAN(4) X MUCK/SILT(2) X TOTAL NUMBER OF SUBSTRATE TYPES: X >4(2) NOTE: (Ignore sludge that originates from point sources: scc COMMENTS:	GRAVEL(7) SAND(6) BEDROCK(5) DETRITUS(3) X ARTIFIC(0) <4(0)	X LIMESTO X TILLS(1) SANDSTO SHALE(-1) COAL FIN	RATE ORIGIN (all) INE(1) RIP/RAP(0) HARDPAN(0) ONE(0)	SILT COV X SILT-HEAVY(-2) SILT-NORM(0)	BSTRATE SCORE 4 (ER (one) SILT-MOD(-1) SILT-FREE(1) edness (check one) MODERATE(-1) NONE(1)	.□
2) INSTREAM COVER: (20) TYPE(Check all UNDERCUT BANKS(1) OVERHANGING VEGETATION(1) X ROOTW X SHALLOWS (IN SLOW WATER)(1) COMMENTS:	OOLS(2) OXBOWS(1 ADS(1) AQUATIC N) IACROPHYTES(1) VOODY DEBRIS(1)	<u>AMOUNT</u>	(Check only one or C EXTENSIVE >76 MODERATE 25- SPARSE 5-25% X NEARLY ABSEN	75%(7) (3)	_
3) CHANNEL MORPHOLOGY: (Check ONLY ONLY ONLY ONLY ONLY ONLY ONLY ONLY	ONE per Category or Ch CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVE	X HIGH(3) MODERA LOW(1)	TY MOS	DIFICATION/OTHI NAGGING ELOCATION ANOPY REMOVAL REDGING NE SIDE CHANNEL MO	IMPOUND ISLAND LEVEED BANK SHAPING	1
L R (per bank) L WIDE>150ft.(4)	R (most predominant property source) FOREST, SWAMP(3) OPEN PASTURE/ROW CRESID., PARK, NEW FIELD FENCED PASTURE(1)	PLAIN QUALITY er bank) L R	GE per bank) (10) (per bank) URBAN OR INDUSTRIA SHURB OR OLD FIELDI CONSERV. TILLAGE(1) MINING/CONSTRUCTIO	L(0) X	RIPARIAN SCORE 5 IK EROSION R (per bank) X NONE OR LITTLE(3) MODERATE(2) HEAVY OR SEVERE(1	•
2.4-4 ft.(4)	<i>'</i>	×	DOL/RUN/RIFFLE CU TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)	JRRENT VELOCITY X EDDIES(1) INTERSTITIAL(- INTERMITTENT		<u> </u>
RIFFLE/RUN DEPTH X GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY 2-4 in.(1) GENERALLY <2 in.(Riffle=0)(0) COMMENTS: 6) GRADIENT (FEET/MILE)(10) 0.4	RIFFLE/RUN SUBSTRATE STABLE (e.g., Cobble,Boul X MOD. STABLE (e.g., Pea G UNSTABLE (Gravel, Sand) NO RIFFLE(0) % POOL 20	der)(2) Gravel)(1)	RIFFLE/RL EXTENS X MODERA LOW(1)	NO NO	•	

STREAM: White River	RIVER MILE	15.9	DATE:	9/20/2004	QHEI SCORE 60.5
1) SUBSTRATE: (Check ONLY Two Substrate POOL RIFFLE BLDER/SLAB(10) BOULDER(9) COBBLE(8) HARDPAN(4) X MUCK/SILT(2) TOTAL NUMBER OF SUBSTRATE TYPES: X >4(2) NOTE: (Ignore sludge that originates from point sources: scc COMMENTS:	POOL GRAVEL(7) SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0) <44(0)	X LIMEST X X TILLS(1 SANDS' SHALE(RATE ORIGIN (all ONE(1) RIP/RAP(0) HARDPAN(SILT COV SILT-HEAVY(-2) SILT-NORM(0)	BSTRATE SCORE 12 /ER (one) SILT-MOD(-1) SILT-FREE(1) edness (check one) X MODERATE(-1) NONE(1)
2) INSTREAM COVER: (20) TYPE(Check all UNDERCUT BANKS(1) X DEEP POOR OVERHANGING VEGETATION(1) ROOTW X SHALLOWS (IN SLOW WATER)(1) BOULDE COMMENTS:	OOLS(2) OXBOWS ADS(1) AQUATIO	S(1) C MACROPHYTES(1) R WOODY DEBRIS(1)	<u>AMOUN'</u>	Check only one or (EXTENSIVE >7: MODERATE 25 SPARSE 5-25% X NEARLY ABSE	75%(7)
3) CHANNEL MORPHOLOGY: (Check ONLY ONLY ONLY ONLY ONLY ONLY ONLY ONLY	DNE per Category or (CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVEN	X HIGH(3) MODER LOW(1)	MATE(2)	ODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MO	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIAN ZONE AND BANK EROSION: (C) River Right Looking Downstream RIPARIAN WIDTH (per bank) L R (per bank) L WIDE>150ft.(4) MODERATE 30-150 ft.(3) X NARROW 15-30 ft.(2) VERY NARROW 3-15 ft.(1) NONE(0) COMMENTS:	R (most predominant FOREST, SWAMP(3) OPEN PASTURE/ROW RESID.,PARK,NEW FIEL FENCED PASTURE(1)	DPLAIN QUALITY per bank) L F		BAI L HAL(0) X D(2)	RIPARIAN SCORE 8.5 NK EROSION R (per bank) X NONE OR LITTLE(3) MODERATE(2) HEAVY OR SEVERE(1)
2.4-4 ft.(4)	′ ′	×	TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)	CURRENT VELOCITY X EDDIES(1) INTERSTITIAL(-	
RIFFLE/RUN DEPTH X GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY 2-4 in.(1) GENERALLY <2 in.(Riffle=0)(0) COMMENTS: 6) GRADIENT (FEET/MILE)(10) 0.4	RIFFLE/RUN SUBSTRA STABLE (e.g., Cobble,Bo X MOD. STABLE (e.g., Pea UNSTABLE (Gravel, Sar NO RIFFLE(0) % POOL 30	oulder)(2) a Gravel)(1)	X MODEI	RATE(0) NO	RIFFLE SCORE 5 E(2) RIFFLE(0) ADJENT SCORE 6

STREAM:	White River		RIVER MILE	24.3	DATE:	9/21/2004	QHEI SCORE 55
TYPE BLDE	ER/SLAB(10)	Two Substrate T	POOL GRAVEL(7) SAND(6) X	X L	JBSTRATE ORIGI	IN (all) SILT CO P/RAP(0) SILT-HEAVY(-2) RDPAN(0) X SILT-NORM(0)	SILT-FREE(1)
X MUC	MBER OF SUBSTRATE TY		BEDROCK(5) DETRITUS(3) X ARTIFIC(0) <4(0) e is based on natural suits.		:ANDSTONE(0) :HALE(-1) :OAL FINES(-2)	Extent of Ember	MODERATE(-1) NONE(1)
UNDERCU'	T BANKS(1) GING VEGETATION(1) S (IN SLOW WATER)(1)	X DEEP POOR ROOTWA	OLS(2) OXBO	WS(1) TIC MACROPHYT OR WOODY DEB	ES(1)	MOUNT (Check only one o EXTENSIVE : MODERATE : SPARSE 5-25 X NEARLY ABS	25-75%(7) 5%(3)
3) CHANNE SINUOSITY HIGH(4) MODERATI X LOW(2) NONE(1) COMMEN	EXCELL GOOD(5 X FAIR(3) POOR(1	DPMENT CENT(7) X	NE per Category of HANNELIZATION NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO REC	ST.	I AVERAGE)(20) ABILITY IIGH(3) IODERATE(2) OW(1)	MODIFICATION/OT SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL IN	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIA	N ZONE AND BAN	· ·	eck ONE box or C	heck 2 and A	/ERAGE per bank	k) (10)	RIPARIAN SCORE 5
RIPARIAN L R (per MOD X X NARF	=>150ft.(4) ERATE 30-150 ft.(3) ROW 15-30 ft.(2) Y NARROW 3-15 ft.(1) E(0)	EROS L F	(Most predominal FOREST, SWAMP(3) OPEN PASTURE/RO RESID.,PARK,NEW F	w CROP(0)	R (per bank) URBAN OR IN SHURB OR OI CONSERV. TII	DUSTRIAL(0) X	ANK EROSION R (per bank) X NONE OR LITTLE(3) MODERATE(2) HEAVY OR SEVERE(1)
•	LIDE AND RIFFLE	•	′	NO POOL = 0			POOL SCORE 10
X >4 ft.(6) 2.4-4 ft.(4) 1.2-2.4 ft.(2) <1.2 ft.(1) <0.6 ft.(Poo	ol=0)(0)	POOL WIE	.OGY (Check 1) DTH>RIFFLE WIDTH(2) DTH=RIFFLE WIDTH(1) DTH <riffle td="" width(0)<=""><td></td><td>TORRENTIAL(X FAST(1) X MODERATE(1) X SLOW(1)</td><td>INTERSTITIA</td><td>L(-1)</td></riffle>		TORRENTIAL(X FAST(1) X MODERATE(1) X SLOW(1)	INTERSTITIA	L(-1)
RIFFLE/RU X GENERALL GENERALL	N DEPTH .Y >4 in. MAX.>20 in.(4) .Y >4 in. MAX.<20 in.(3) .Y 2-4 in.(1) .Y <2 in.(Riffle=0)(0)		STABLE (e.g., Cobble MOD. STABLE (e.g., UNSTABLE (Gravel, S NO RIFFLE(0)	e,Boulder)(2) Pea Gravel)(1)			RIFFLE SCORE 5 SS DNE(2) NO RIFFLE(0)
6) GRADIE	NT (FEET/MILE)(10	0.4	% POOL3	0 % RI	FFLE 15	%RUN_ 55 G	RADIENT SCORE 6

STREAM:	White River	RIVER MILE	39.2	DATE:	9/22/2004	QHEI SCORE 47.5
1) SUBSTR	ATE: (Check ONLY Two S	Substrate Type Boxes: Check a		t)(20) TRATE ORIGIN (all		BSTRATE SCORE 6 /ER (one)
BOUL COBE HARE X MUCH TOTAL NUM		GRAVEL(7) X SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0) >4(2) X <4(0) sources: score is based on natural substr	TILLS SAND SHAL COAL	STONE(0)	(0) SILT-NORM(0)	X SILT-MOD(-1) SILT-FREE(1) ledness (check one) X MODERATE(-1) NONE(1)
2) INSTREA	AM COVER: (20)	Check all that apply)		AMOUN	T (Check only one or	COVER SCORE 5 Check 2 and AVERAGE)
OVERHANO	T BANKS(1) GING VEGETATION(1) S (IN SLOW WATER)(1)	X DEEP POOLS(2) OXBOWS ROOTWADS(1) AQUATIC	(1) MACROPHYTES(1 WOODY DEBRIS()	EXTENSIVE >7 MODERATE 25 SPARSE 5-25% X NEARLY ABSE	5%(11) -75%(7) (3)
3) CHANNE SINUOSITY HIGH(4) MODERATE X LOW(2) NONE(1)	DEVELOPMEN EXCELLENT(7)	CONLY ONE per Category or CONT CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVERED	STABI HIGH X MODE LOW(LITY M 3) ERATE(2)	ODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MC	IMPOUND ISLAND LEVEED BANK SHAPING
COMMENT		SION: (Check ONE box or Che	ck 2 and AVER	PAGE ner hank) (10)	RIPARIAN SCORE 5.5
River Right RIPARIAN L R (per WIDE MODI X X NARR	Looking Downstream WIDTH (per bank) bank) >>150ft.(4) ERATE 30-150 ft.(3) ROW 15-30 ft.(2) V NARROW 3-15 ft.(1) E(0)	EROSION/RUNOFF-FLOOI L R (most predominant FOREST, SWAMP(3) OPEN PASTURE/ROW O RESID.,PARK,NEW FIEL FENCED PASTURE(1)	DPLAIN QUALIT		BA L RIAL(0) X LD(2) X	NK EROSION R (per bank) X NONE OR LITTLE(3) MODERATE(2) HEAVY OR SEVERE(1)
,) -0)(0)	MORPHOLOGY (Check 1) X POOL WIDTH>RIFFLE WIDTH(2) POOL WIDTH POOL WIDTH POOL WIDTH RIFFLE WIDTH(0)		POOL/RUN/RIFFLE TORRENTIAL(-1) X FAST(1) X MODERATE(1) X SLOW(1)	CURRENT VELOCITY X EDDIES(1) INTERSTITIAL(INTERMITTEN	
GENERALL GENERALL GENERALL COMMEN	Y >4 in. MAX.>20 in.(4) Y >4 in. MAX.<20 in.(3) Y 2-4 in.(1) Y <2 in.(Riffle=0)(0)	RIFFLE/RUN SUBSTRA' STABLE (e.g., Cobble,Bo X MOD. STABLE (e.g., Pea UNSTABLE (Gravel, San	ulder)(2) Gravel)(1) d)(0)	X MODE	RATE(0) NC	IE(2) RIFFLE(0)
6) GRADIEI	NT (FEET/MILE)(10) (0.8 % POOL 1	% RIFFL	.E 0 %	RUN 99 GR	ADIENT SCORE 8

STREAM: White River	RIVER MILE	45.7	DATE:	9/23/2004	QHEI SCORE 44
1) SUBSTRATE: (Check ONLY Two Substrate POOL RIFFLE BLDER/SLAB(10) BOULDER(9) COBBLE(8) X HARDPAN(4) X X MUCK/SILT(2) TOTAL NUMBER OF SUBSTRATE TYPES: X >4(2) NOTE: (Ignore sludge that originates from point sources: scc	POOL GRAVEL(7) SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0) 44(0)	X LIMES X LIMES TILLS(SANDS SHALE COAL	TRATE ORIGIN (all TONE(1) RIP/RAP(0 HARDPAN(0)	SILT CON SILT-HEAVY(-2) SILT-NORM(0)	BSTRATE SCORE 4 /ER (one) X SILT-MOD(-1) SILT-FREE(1) ledness (check one) X MODERATE(-1) NONE(1)
2) INSTREAM COVER: (20) TYPE(Check all UNDERCUT BANKS(1) OVERHANGING VEGETATION(1) X SHALLOWS (IN SLOW WATER)(1) COMMENTS:	OOLS(2) OXBOW ADS(1) AQUATI	S(1) C MACROPHYTES(1) R WOODY DEBRIS(1		T (Check only one or of extensive >7 MODERATE 25 SPARSE 5-25% X NEARLY ABSE	-75%(7) -(3)
3) CHANNEL MORPHOLOGY: (Check ONLY ONLY OF SINUOSITY DEVELOPMENT) HIGH(4) EXCELLENT(7) GOOD(5) LOW(2) FAIR(3) X NONE(1) X POOR(1) COMMENTS:	CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVERING	X HIGH(LITY M B) RATE(2)	ODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MC	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIAN ZONE AND BANK EROSION: (C River Right Looking Downstream	heck ONE box or Ch SION/RUNOFF-FLOC R (most predominan FOREST, SWAMP(3) OPEN PASTURE/ROW RESID.,PARK,NEW FIE FENCED PASTURE(1)	DDPLAIN QUALIT t per bank) L CROP(0)		BAI L MIAL(0) X D(2)	RIPARIAN SCORE 6 NK EROSION R (per bank) X NONE OR LITTLE(3) MODERATE(2) HEAVY OR SEVERE(1)
X >4 ft.(6) POOL W 2.4-4 ft.(4) POOL W	DLOGY (Check 1) IDTH>RIFFLE WIDTH(2) IDTH=RIFFLE WIDTH(1) IDTH <riffle td="" width(0)<=""><td></td><td>POOL/RUN/RIFFLE of torrential(-1) X FAST(1) X MODERATE(1) X SLOW(1)</td><td>X EDDIES(1) INTERSTITIAL(</td><td></td></riffle>		POOL/RUN/RIFFLE of torrential(-1) X FAST(1) X MODERATE(1) X SLOW(1)	X EDDIES(1) INTERSTITIAL(
GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY 2-4 in.(1) GENERALLY <2 in.(Riffle=0)(0) COMMENTS: 6) GRADIENT (FEET/MILE)(10) 0.9	STABLE (e.g., Cobble,E MOD. STABLE (e.g., Pe UNSTABLE (Gravel, Sa X NO RIFFLE(0)	Boulder)(2) ea Gravel)(1)	EXTEN MODE LOW(1	RATE(0) NO	RIFFLE SCORE 0

STREAM: WFWR	RIVER MILE	57.5	DATE:	9/21/2004	QHEI SCORE 46
1) SUBSTRATE: (Check ONLY Two Substrate Type POOL RIFFLE BLDER/SLAB(10)	/Pe Boxes: Check all POOL RI GRAVEL(7) SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0)	types present)(20 FFLE SUBSTRA LIMESTON X TILLS(1) SANDSTO SHALE(-1) COAL FINE	D) ATE ORIGIN (all) IE(1) RIP/RAP(0) HARDPAN(0) NE(0)	SILT CO SILT-HEAVY(-2) X SILT-NORM(0)	UBSTRATE SCORE 10 VER (one) SILT-MOD(-1) SILT-FREE(1) dedness (check one) X MODERATE(-1) NONE(1)
2) INSTREAM COVER: (20) TYPE(Check all th UNDERCUT BANKS(1) OVERHANGING VEGETATION(1) X ROOTWAI X SHALLOWS (IN SLOW WATER)(1) COMMENTS:	OLS(2) OXBOWS(1) ACROPHYTES(1) VOODY DEBRIS(1)	<u>AMOUNT</u> ((Check only one or EXTENSIVE >7 X MODERATE 25 SPARSE 5-259 NEARLY ABSE	5-75%(7) 6(3)
	IE per Category or Ch HANNELIZATION NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVE	STABILITY HIGH(3) MODERAT X LOW(1)	Y MOE SN E(2) RE CA	DIFICATION/OTH LAGGING ELOCATION LNOPY REMOVAL REDGING NE SIDE CHANNEL MO	IMPOUND ISLAND LEVEED BANK SHAPING
L R (per bank) L R WIDE>150ft.(4) X	iON/RUNOFF-FLOODI (most predominant purification) FOREST, SWAMP(3) OPEN PASTURE/ROW CR RESID.,PARK,NEW FIELD FENCED PASTURE(1)	PLAIN QUALITY er bank) L R (E per bank) (10) per bank) JRBAN OR INDUSTRIAL SHURB OR OLD FIELD(2 CONSERV. TILLAGE(1) JINING/CONSTRUCTION	L (0) 2) X	RIPARIAN SCORE 6 NK EROSION R (per bank) NONE OR LITTLE(3) MODERATE(2) X HEAVY OR SEVERE(1)
>4 ft.(6) POOL WIE 2.4-4 ft.(4)	2) NO F OGY (Check 1) TH>RIFFLE WIDTH(2) TH=RIFFLE WIDTH(1) TH <riffle td="" width(0)<=""><td>I T</td><td>OL/RUN/RIFFLE CU ORRENTIAL(-1) :AST(1) MODERATE(1) SLOW(1)</td><td>RRENT VELOCITY EDDIES(1) INTERSTITIAL INTERMITTEN</td><td>Τ(-2)</td></riffle>	I T	OL/RUN/RIFFLE CU ORRENTIAL(-1) :AST(1) MODERATE(1) SLOW(1)	RRENT VELOCITY EDDIES(1) INTERSTITIAL INTERMITTEN	Τ(-2)
GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY 2-4 in.(1) GENERALLY <2 in.(Riffle=0)(0) COMMENTS:	FFLE/RUN SUBSTRATE STABLE (e.g., Cobble,Boul MOD. STABLE (e.g., Pea G UNSTABLE (Gravel, Sand) NO RIFFLE(0)	der)(2) iravel)(1)	RIFFLE/RUI EXTENSIV MODERAT LOW(1)	TE(0) X NO	RIFFLE SCORE 0 S NE(2) D RIFFLE(0) RADIENT SCORE 8

STREAM: WFWR	RIVER MILE	61.52 DATE	E: <u>9/21/2004</u>	QHEI SCORE 53
1) SUBSTRATE: (Check ONLY Two Substance of Substrate types:	X	RIFFLE SUBSTRATE OR LIMESTONE(1) X TILLS(1) SANDSTONE(0) SHALE(-1) COAL FINES(-2)	RIP/RAP(0) SILT-HEAVY(-2) HARDPAN(0) X SILT-NORM(0)	UBSTRATE SCORE 12 OVER (one) SILT-MOD(-1) SILT-FREE(1) Idedness (check one) MODERATE(-1) NONE(1)
UNDERCUT BANKS(1) X OVERHANGING VEGETATION(1) X RC	_	(1) MACROPHYTES(1) WOODY DEBRIS(1)	AMOUNT (Check only one of EXTENSIVE > X MODERATE 2 SPARSE 5-25 NEARLY ABS	-75%(11) 25-75%(7) %(3)
3) CHANNEL MORPHOLOGY: (Check ON SINUOSITY DEVELOPMENT HIGH(4) EXCELLENT(7) GOOD(5) X LOW(2) X FAIR(3) NONE(1) POOR(1) COMMENTS:	LY ONE per Category or C CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVERING OF	STABILITY HIGH(3) MODERATE(2) X LOW(1)	MODIFICATION/OTE SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL M	IMPOUND ISLAND LEVEED BANK SHAPING
L R (per bank)	EROSION/RUNOFF-FLOOI L R (most predominant X FOREST, SWAMP(3) X OPEN PASTURE/ROW (C) RESID.,PARK,NEW FIEL FENCED PASTURE(1)	DPLAIN QUALITY per bank) L R (per bar URBAN OF CROP(0) SHURB OF CONSERV	<u>B</u>	RIPARIAN SCORE 7 ANK EROSION R (per bank) NONE OR LITTLE(3) MODERATE(2) X HEAVY OR SEVERE(1)
>4 ft.(6) PC PC PC	NO RPHOLOGY (Check 1) OL WIDTH>RIFFLE WIDTH(2) OL WIDTH=RIFFLE WIDTH(1) OL WIDTH <riffle td="" width(0)<=""><td>POOL = 0 POOL/RUN TORRENT FAST(1) MODERAT SLOW(1)</td><td>INTERSTITIAL</td><td>L(-1) NT(-2)</td></riffle>	POOL = 0 POOL/RUN TORRENT FAST(1) MODERAT SLOW(1)	INTERSTITIAL	L(-1) NT(-2)
GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.>20 in.(3) GENERALLY >4 in. MAX. GENERALLY >4 in. (1) X GENERALLY <2 in.(Riffle=0)(0) COMMENTS: 6) GRADIENT (FEET/MILE)(10) 1.02	RIFFLE/RUN SUBSTRA STABLE (e.g., Cobble, Bo MOD. STABLE (e.g., Pea UNSTABLE (Gravel, Sand X NO RIFFLE(0)	ulder)(2) Gravel)(1) d)(0)	MODERATE(0) X N	RIFFLE SCORE 0 SS DNE(2) O RIFFLE(0) RADIENT SCORE 8
6) GRADIENT (FEET/MILE)(10) 1.02	% POOL	% RIFFLE	% RUN 100 G	KADIENT SCORE O

STREAM:	WFWR	RIVER MILE	73.06	DATE:	9/22/2004	QHEI SCORE 62
TYPE	POOL RIFFL	Substrate Type Boxes: Check	RIFFLE SUBS	TRATE ORIGIN (a	all) SILT CC	UBSTRATE SCORE 13
X COBI HARI X MUC TOTAL NUM NOTE: (Igno	DPAN(4) K/SILT(2) IBER OF SUBSTRATE TYPES: re sludge that originates from poin	X GRAVEL(7) X SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0) >4(2) X <4(0) at sources: score is based on natural subst	X TILLS(SAND: SHALE	STONE(0)	N(0) X SILT-NORM(0)	SILT-MOD(-1) SILT-FREE(1) Idedness (check one) X MODERATE(-1) NONE(1)
2) INSTREA	AM COVER: (20)					COVER SCORE 16
OVERHAN	T BANKS(1) GING VEGETATION(1) S (IN SLOW WATER)(1)		S(1) C MACROPHYTES(1) R WOODY DEBRIS(1		NT (Check only one or X EXTENSIVE > MODERATE 2 SPARSE 5-25 NEARLY ABSI	25-75%(7) %(3)
3) CHANNE	EL MORPHOLOGY: (Che	ck ONLY ONE per Category or		,, ,		16
X HIGH(4) MODERAT LOW(2) NONE(1)	EXCELLENT(7)	NT CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECO	X LOW(1	3) RATE(2)	MODIFICATION/OTI SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL M	IMPOUND ISLAND LEVEED BANK SHAPING
COMMEN	TS:					
River Right	N ZONE AND BANK ERO Looking Downstream WIDTH (per bank)	OSION: (Check ONE box or Che EROSION/RUNOFF-FLOC				RIPARIAN SCORE 7 ANK EROSION
X WIDE X MOD		L R (most predominant X FOREST, SWAMP(3) X OPEN PASTURE/ROW RESID.,PARK,NEW FIE FENCED PASTURE(1)	CROP(0)	R (per bank) URBAN OR INDUS' SHURB OR OLD FI CONSERV. TILLAG MINING/CONSTRU	ELD(2) X EE(1)	R (per bank) NONE OR LITTLE(3) MODERATE(2) X HEAVY OR SEVERE(1)
MAX. DEF >4 ft.(6) 2.4-4 ft.(4) 1.2-2.4 ft.(2) <1.2 ft.(1) <0.6 ft.(Poc		QUALITY (12) MORPHOLOGY (Check 1) POOL WIDTH>RIFFLE WIDTH(2) POOL WIDTH+RIFFLE WIDTH(1) POOL WIDTH+RIFFLE WIDTH(0)	D POOL = 0	POOL/RUN/RIFFLE TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)	E CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN	
GENERALI GENERALI	_Y >4 in. MAX.>20 in.(4) _Y >4 in. MAX.<20 in.(3) _Y 2-4 in.(1) _Y <2 in.(Riffle=0)(0)	RIFFLE/RUN SUBSTRA STABLE (e.g., Cobble,B MOD. STABLE (e.g., Pe UNSTABLE (Gravel, Sai X NO RIFFLE(0)	oulder)(2) a Gravel)(1)	EXT	DERATE(0) X N	RIFFLE SCORE 0 SS ONE(2) O RIFFLE(0)
6) GRADIE	NT (FEET/MILE)(10)	1.28 % POOL	% RIFFL	E %	% RUN 100 gi	RADIENT SCORE 10

STREAM: WFWR		RIVER MILE	84.5	DATE:	10/19/2004	QHEI SCORE 53
TYPE BLDER/SLAB(10) BOULDER(9) COBBLE(8) HARDPAN(4) X MUCK/SILT(2) TOTAL NUMBER OF SUBS		POOL GRAVEL(7) X SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0) X <4(0)	RIFFLE SUBS LIME X TILLS SANG SHAL COAL	STRATE ORIGIN STONE(1) RIP/R	AP(0) X SILT-NORM(0)	UBSTRATE SCORE 12 VER (one) SILT-MOD(-1) SILT-FREE(1) Idedness (check one) MODERATE(-1) NONE(1)
2) INSTREAM COVER UNDERCUT BANKS(1) X OVERHANGING VEGETA: X SHALLOWS (IN SLOW WA	TYPE(Check all to DEEP PO	OOLS(2) OXBOW ADS(1) AQUATI	S(1) C MACROPHYTES(* R WOODY DEBRIS(1)	UNT (Check only one or EXTENSIVE > X MODERATE 2 SPARSE 5-25 NEARLY ABSI	5-75%(7) %(3)
*	EXCELLENT(7) GOOD(5)	CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVERED	STAB HIGH X MOD LOW	ILITY I(3) ERATE(2)	MODIFICATION/OTI SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL M	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIAN ZONE AN River Right Looking Do RIPARIAN WIDTH (pei L R (per bank) X WIDE>150ft.(4) MODERATE 30-150 NARROW 15-30 ft.(2) VERY NARROW 3-1 NONE(0) COMMENTS:	ft.(3) ERO L X ft.(3)	Neck ONE box or Ch SION/RUNOFF-FLOC R (most predominan X FOREST, SWAMP(3) OPEN PASTURE/ROW RESID.,PARK,NEW FIE FENCED PASTURE(1)	DDPLAIN QUALI t per bank) L CROP(0)		B/L L X STRIAL(0) X FIELD(2) AGE(1)	RIPARIAN SCORE 9 ANK EROSION R (per bank) NONE OR LITTLE(3) X MODERATE(2) HEAVY OR SEVERE(1)
5) POOL/GLIDE AND F MAX. DEPTH (Check 24 ft.(6) 2.4-4 ft.(4) 1.2-2.4 ft.(2) <1.2 ft.(1) <0.6 ft.(Pool=0)(0) COMMENTS: No po	POOL W	12) NOLOGY (Check 1) IDTH>RIFFLE WIDTH(2) IDTH=RIFFLE WIDTH(1) IDTH <riffle td="" width(0)<=""><td>O POOL = 0</td><td>POOL/RUN/RIFF TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)</td><td>LE CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN</td><td>.(-1)</td></riffle>	O POOL = 0	POOL/RUN/RIFF TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)	LE CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN	.(-1)
RIFFLE/RUN DEPTH GENERALLY >4 in. MAX.> GENERALLY >4 in. MAX.< GENERALLY 2-4 in.(1) GENERALLY <2 in.(Riffle= COMMENTS:	20 in.(4) 20 in.(3) 0)(0)	RIFFLE/RUN SUBSTR/ STABLE (e.g., Cobble,E MOD. STABLE (e.g., Pe UNSTABLE (Gravel, Sa X NO RIFFLE(0)	Boulder)(2) ea Gravel)(1)	Ex MC	DDERATE(0) X N	RIFFLE SCORE 0 SS INE(2) O RIFFLE(0)

STREAM:	WFWR	RIVER MILE	101.4	DATE:	9/20/2004	QHEI SCORE 57
1) SUBSTR	RATE: (Check ONLY Two S	Substrate Type Boxes: Check a		:)(20) FRATE ORIGIN (a		UBSTRATE SCORE 16
BOUI COBI HARI MUCI TOTAL NUM		X X GRAVEL(7) X SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0) >4(2) X 34(0) sources: score is based on natural substi	X TILLS(SAND: SHALE	STONE(0)	N(0) X SILT-NORM(0)	SILT-MOD(-1) SILT-FREE(1) idedness (check one) MODERATE(-1) X NONE(1)
_	AM COVER: (20) <u>TYPE(</u> (T BANKS(1)	Check all that apply)	2/4)	AMOU	NT (Check only one or	COVER SCORE 10 Check 2 and AVERAGE)
X OVERHAN	GING VEGETATION(1) S (IN SLOW WATER)(1)	ROOTWADS(1) AQUATIO	C MACROPHYTES(1) R WOODY DEBRIS(1)		X MODERATE 2 SPARSE 5-25 NEARLY ABSI	25-75%(7) %(3)
3) CHANNE SINUOSITY HIGH(4) X MODERATI LOW(2) NONE(1)	Y DEVELOPMEN EXCELLENT(7)	IT CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVERED	STABII HIGH(MODE X LOW(LITY 3) RATE(2)	MODIFICATION/OTI SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL M	IMPOUND ISLAND LEVEED BANK SHAPING
COMMEN	TS:					
River Right RIPARIAN L R (per X WIDE MOD NARR	Looking Downstream WIDTH (per bank) bank) =>150ft.(4) ERATE 30-150 ft.(3) ROW 15-30 ft.(2) Y NARROW 3-15 ft.(1) E(0)	EROSION/RUNOFF-FLOO L R (most predominant X FOREST, SWAMP(3) OPEN PASTURE/ROW RESID.,PARK,NEW FIEL FENCED PASTURE(1)	DPLAIN QUALIT per bank) L CROP(0)		BA L TRIAL(0) ELD(2) E(1) X	RIPARIAN SCORE 6 ANK EROSION R (per bank) NONE OR LITTLE(3) X MODERATE(2) HEAVY OR SEVERE(1)
MAX. DEF >4 ft.(6) 2.4-4 ft.(4) 1.2-2.4 ft.(2) <1.2 ft.(1) <0.6 ft.(Pool	•	MORPHOLOGY (Check 1) POOL WIDTH>RIFFLE WIDTH(2) POOL WIDTH=RIFFLE WIDTH(1) POOL WIDTH <riffle td="" width(0)<=""><td>POOL = 0</td><td>POOL/RUN/RIFFLE TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)</td><td>E CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN</td><td></td></riffle>	POOL = 0	POOL/RUN/RIFFLE TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)	E CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN	
GENERALL GENERALL	.Y >4 in. MAX.>20 in.(4) .Y >4 in. MAX.<20 in.(3) .Y >2-4 in.(1) .Y <2 in.(Riffle=0)(0) TS:	RIFFLE/RUN SUBSTRA STABLE (e.g., Cobble,Bo MOD. STABLE (e.g., Pea UNSTABLE (Gravel, Sar X NO RIFFLE(0)	oulder)(2) a Gravel)(1) ad)(0)	EXTI MOD	X N (1)	ONE(2) O RIFFLE(0)
6) GRADIE	NT (FEET/MILE)(10) 1	1.20 % POOL	% RIFFL	E 9	6 RUN 100 gi	RADIENT SCORE 10

STREAM: West Fork White River	RIVER MILE	105.7	DATE:	9/20/2004	QHEI SCORE 56.5
1) SUBSTRATE: (Check ONLY Two Substrater Type POOL RIFFLE BLDER/SLAB(10) X COBBLE(8)	GRAVEL(7) X SAND(6) X BEDROCK(5) DETRITUS(3) ARTIFIC(0) X	all types present) RIFFLE SUBST LIMEST X TILLS(1 SANDS SHALE(COAL F	(20) RATE ORIGIN (all) ONE(1) RIP/RAP(0) HARDPAN(0) TONE(0)	SILT COV	BSTRATE SCORE 13.5 VER (one) X SILT-MOD(-1) SILT-FREE(1) dedness (check one) X MODERATE(-1) NONE(1)
UNDERCUT BANKS(1) OVERHANGING VEGETATION(1) X DEE X POO	all that apply) EP POOLS(2) OTWADS(1) OXBOWS		AMOUNT	EXTENSIVE >7 X MODERATE 25 SPARSE 5-25% NEARLY ABSE	5-75%(7) 5-(3)
3) CHANNEL MORPHOLOGY: (Check ONL SINUOSITY DEVELOPMENT HIGH(4) EXCELLENT(7) MODERATE(3) GOOD(5) X LOW(2) FAIR(3) NONE(1) X POOR(1) COMMENTS:	Y ONE per Category or C CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECO	STABILI HIGH(3) MODER X LOW(1)	ATTY MC	DDIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MO	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIAN ZONE AND BANK EROSION River Right Looking Downstream RIPARIAN WIDTH (per bank) L R (per bank) UNIDE>150ft.(4) X MODERATE 30-150 ft.(3) NARROW 15-30 ft.(2) VERY NARROW 3-15 ft.(1) NONE(0) COMMENTS:	ROSION/RUNOFF-FLOO R (most predominant	DPLAIN QUALITY per bank) L F		AL(0) X D(2) X	NK EROSION R (per bank) NONE OR LITTLE(3) X MODERATE(2) HEAVY OR SEVERE(1)
X >4 ft.(6) X POO 2.4-4 ft.(4) POO	PHOLOGY (Check 1) DL WIDTH>RIFFLE WIDTH(2) DL WIDTH=RIFFLE WIDTH(1) DL WIDTH <riffle td="" width(0)<=""><td></td><td>TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)</td><td>X EDDIES(1) INTERSTITIAL</td><td></td></riffle>		TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)	X EDDIES(1) INTERSTITIAL	
RIFFLE/RUN DEPTH GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY >4 in.(1) GENERALLY <2 in.(Riffle=0)(0) COMMENTS: 6) GRADIENT (FEET/MILE)(10) 0.93	RIFFLE/RUN SUBSTRA STABLE (e.g., Cobble,B MOD. STABLE (e.g., Per UNSTABLE (Gravel, Sar NO RIFFLE(0) % POOL 25	oulder)(2) a Gravel)(1) nd)(0)	EXTENS MODER LOW(1)	PATE(0) NC	RIFFLE SCORE 0 S NE(2) PRIFFLE(0) ADJENT SCORE 6

STREAM: West Fork White River	RIVER MILE	115.5	DATE:	9/23/2004	QHEI SCORE 44.5
1) SUBSTRATE: (Check ONLY Two Subs	trate Type Boxes: Check POOL X GRAVEL(7) X SAND(6) BEDROCK(5) DETRITUS(3) ARTIFIC(0) X <4(0)	all types present)(RUN SUBSTI X LIMESTI X X TILLS(1) SANDSTI SHALE(- COAL FI	20) RATE ORIGIN (al DNE(1) RIP/RAP(I HARDPAN TONE(0)	SILT CO SILT CO SILT-HEAVY(-2) SILT-NORM(0)	JBSTRATE SCORE 11.5 VER (one) X SILT-MOD(-1) SILT-FREE(1) dedness (check one) X MODERATE(-1) NONE(1)
X UNDERCUT BANKS(1) OVERHANGING VEGETATION(1) X R	—	(S(1) C MACROPHYTES(1) OR WOODY DEBRIS(1)	AMOUN	T (Check only one or EXTENSIVE >7 MODERATE 26 X SPARSE 5-259 NEARLY ABSE	5-75%(7) 6(3)
3) CHANNEL MORPHOLOGY: (Check ON SINUOSITY DEVELOPMENT HIGH(4) EXCELLENT(7) MODERATE(3) GOOD(5) LOW(2) FAIR(3) X NONE(1) X POOR(1) COMMENTS:	ILY ONE per Category or CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECOVERING OF	STABILI HIGH(3) X MODER LOW(1)	TY <u>M</u>	ODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MO	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIAN ZONE AND BANK EROSIO River Right Looking Downstream RIPARIAN WIDTH (per bank) L R (per bank) WIDE>150ft.(4) X X MODERATE 30-150 ft.(3) NARROW 15-30 ft.(2) VERY NARROW 3-15 ft.(1) NONE(0) COMMENTS:	N: (Check ONE box or Ch EROSION/RUNOFF-FLOC L R (most predominan FOREST, SWAMP(3) X OPEN PASTURE/ROW RESID.,PARK,NEW FIE FENCED PASTURE(1)	DDPLAIN QUALITY t per bank) L R CROP(0)		BA L X X LD(2) (1)	RIPARIAN SCORE 6 NK EROSION R (per bank) X NONE OR LITTLE(3) MODERATE(2) HEAVY OR SEVERE(1)
>4 ft.(6) P(ITY (12) RPHOLOGY (Check 1) DOL WIDTH>RIFFLE WIDTH(2) DOL WIDTH=RIFFLE WIDTH(1) DOL WIDTH <riffle td="" width(0)<=""><td>O POOL = 0</td><td>OOL/RUN/RIFFLE TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)</td><td>CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN</td><td></td></riffle>	O POOL = 0	OOL/RUN/RIFFLE TORRENTIAL(-1) FAST(1) MODERATE(1) SLOW(1)	CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN	
RIFFLE/RUN DEPTH X GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY 2-4 in.(1) GENERALLY <2 in.(Riffle=0)(0) COMMENTS: 6) GRADIENT (FEET/MILE)(10) 0.93	RIFFLE/RUN SUBSTRA STABLE (e.g., Cobble,E MOD. STABLE (e.g., Pe X UNSTABLE (Gravel, Sa NO RIFFLE(0)	Soulder)(2) ea Gravel)(1) and)(0)	X LOW(RATE(0) NO	RIFFLE SCORE 5 S NE(2) PRIFFLE(0) RADIENT SCORE 6

STREAM:	West Fork Wh	ite River	RIVER MILE	125.0	DATE:	9/21/2004	QHEI SCORE 60.5
TYPE BLDE BOUI COBI HARRI MUC TOTAL NUM	ATE: (Check ONLY POOL R/SLAB(10) DER(9) BLE(8) DPAN(4) K/SLT(2) X BBER OF SUBSTRATE TY re sludge that originates fro TS:	X X X X X X X X X X	POOL X GRAVEL(7) X X SAND(6) X BEDROCK(5) DETRITUS(3) X ARTIFIC(0) 44(0)	X Lin X X TIL SA SH	SSTRATE ORIGIN MESTONE(1) X RIP/RA	(all) SILT CO AP(0) SILT-HEAVY(-2) PAN(0) X SILT-NORM(0)	JBSTRATE SCORE 14 VER (one) SILT-MOD(-1) SILT-FREE(1) dedness (check one) X MODERATE(-1) NONE(1)
UNDERCU OVERHAN	T BANKS(1) GING VEGETATION(1) S (IN SLOW WATER)(1)	YPE(Check all to X DEEP PC X ROOTW/BOULDE	OOLS(2) OXBO	WS(1) TIC MACROPHYTE: OR WOODY DEBR	S(1)	UNT (Check only one or EXTENSIVE > X MODERATE 2 SPARSE 5-25' NEARLY ABSE	5-75%(7) %(3)
3) CHANNE SINUOSITY HIGH(4) MODERAT X LOW(2) NONE(1) COMMEN	EXCELLE GOOD(5) X FAIR(3) POOR(1)	PMENT (7) ENT(7)	NE per Category of CHANNELIZATION NONE(6) RECOVERED(4) X RECOVERING(3) X RECENT OR NO REC	STA HIG X MG	AVERAGE)(20) BILITY GH(3) DERATE(2) W(1)	MODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MI	IMPOUND ISLAND LEVEED BANK SHAPING
River Right RIPARIAN L R (per MOD X X NARI	E>150ft.(4) ERATE 30-150 ft.(3) ROW 15-30 ft.(2) (NARROW 3-15 ft.(1) E(0)	m <u>ERO</u> L	SION/RUNOFF-FLC R (most predomina FOREST, SWAMP(3) X OPEN PASTURE/RO RESID.,PARK,NEW F FENCED PASTURE(DODPLAIN QUA	LITY	BA L STRIAL(0) STELD(2) X (GE(1)	RIPARIAN SCORE 4.5 NK EROSION R (per bank) NONE OR LITTLE(3) X MODERATE(2) HEAVY OR SEVERE(1)
•	i=0)(0)	MORPHO POOL WI	12) LOGY (Check 1) IDTH>RIFFLE WIDTH(2 IDTH=RIFFLE WIDTH(1 IDTH <riffle td="" width(0)<=""><td>)</td><td>POOL/RUN/RIFFI TORRENTIAL(-1) X FAST(1) X MODERATE(1) X SLOW(1)</td><td>LE CURRENT VELOCIT X EDDIES(1) INTERSTITIAL INTERMITTEN</td><td>(-1)</td></riffle>)	POOL/RUN/RIFFI TORRENTIAL(-1) X FAST(1) X MODERATE(1) X SLOW(1)	LE CURRENT VELOCIT X EDDIES(1) INTERSTITIAL INTERMITTEN	(-1)
GENERALL GENERALL GENERALL COMMEN	Y >4 in. MAX.>20 in.(4) Y >4 in. MAX.<20 in.(3) Y 2-4 in.(1) Y <2 in.(Riffle=0)(0)	2	STABLE (e.g., Cobble MOD. STABLE (e.g., X UNSTABLE (Gravel, NO RIFFLE(0)	e,Boulder)(2) Pea Gravel)(1)	X MC	DDERATE(0) NO	RIFFLE SCORE 4 S NE(2) D RIFFLE(0)

STREAM:	West Fork Wh	ite River	RIVER MILE	134.7	DATE:	9/21/2004	QHEI SCORE	61.5
TYPE	RATE: (Check ONLY POOL Rr/SLAB(10)	RIFFLE X X	POOL GRAVEL(7) X	RIFFLE SUBS	STRATE ORIGIN		JBSTRATE SCORE VER (one) SILT-MOD(-1)	12
COBE HARE MUCI	DER(9) 3LE(8) DPAN(4) K/SILT(2) IBER OF SUBSTRATE TYI re sludge that originates fro	X X X	BEDROCK(5) DETRITUS(3) ARTIFIC(0) 44(0) e is based on natural subs	SHAI	S(1) HARD DSTONE(0) LE(-1) L FINES(-2)	PAN(0) X SILT-NORM(0) Extent of Embed X EXTENSIVE(-2) LOW(0)	SILT-FREE(1) dedness (check one MODERATE(-1) NONE(1)	<u>3)</u>
X UNDERCU		YPE(Check all the DEEP POINT ROOTWA BOULDER	OLS(2) OXBOW AQUATI	'S(1) C MACROPHYTES(IR WOODY DEBRIS	1)	OUNT (Check only one or X EXTENSIVE > MODERATE 2 SPARSE 5-256	75%(11) 5-75%(7) %(3)	
COMMEN	TS:					NEARLY ABSE	:N1 <5%(1)	
3) CHANNE SINUOSITY HIGH(4) MODERATE X LOW(2) NONE(1)	EXCELLE	PMENT C	NE per Category or CHANNELIZATION NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECO	STAB X HIGH MOD LOW	H(3) DERATE(2)	MODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL M	IMPOUND ISLAND LEVEED BANK SHAPING	12
COMMEN	TS:							
River Right RIPARIAN L R (per X WIDE MODE X NARR	E>150ft.(4) ERATE 30-150 ft.(3) ROW 15-30 ft.(2) / NARROW 3-15 ft.(1) E(0)	m <u>EROS</u> L F	eck ONE box or Ch SION/RUNOFF-FLOC (most predominan FOREST, SWAMP(3) OPEN PASTURE/ROW RESID.,PARK,NEW FIE FENCED PASTURE(1)	DDPLAIN QUALI t per bank) L CROP(0) ELD(1)		BA L JSTRIAL(0) FIELD(2) AGE(1) X	RIPARIAN SCORE NK EROSION R (per bank) NONE OR LITTLE(: X MODERATE(2) HEAVY OR SEVER	(3)
•	LIDE AND RIFFLE/F	,	· —	O POOL = 0			POOL SCORE	
X >4 ft.(6) 2.4-4 ft.(4) 1.2-2.4 ft.(2) <1.2 ft.(1) <0.6 ft.(Poo	I=0)(0)	X POOL WIE	.OGY (Check 1) DTH>RIFFLE WIDTH(2) DTH=RIFFLE WIDTH(1) DTH <riffle td="" width(0)<=""><td></td><td>TORRENTIAL(-1 FAST(1) MODERATE(1) X SLOW(1)</td><td>ELE CURRENT VELOCIT X EDDIES(1) INTERSTITIAL INTERMITTEN</td><td>(-1)</td><td><u>V)</u></td></riffle>		TORRENTIAL(-1 FAST(1) MODERATE(1) X SLOW(1)	ELE CURRENT VELOCIT X EDDIES(1) INTERSTITIAL INTERMITTEN	(-1)	<u>V)</u>
GENERALL GENERALL	Y >4 in. MAX.>20 in.(4) Y >4 in. MAX.<20 in.(3) Y 2-4 in.(1) Y <2 in.(Riffle=0)(0)	<u> </u>	STABLE (e.g., Cobble,E MOD. STABLE (e.g., Pe UNSTABLE (Gravel, Sa NO RIFFLE(0)	Boulder)(2) ea Gravel)(1)	E:		RIFFLE SCORE S NE(2) O RIFFLE(0)	0
6) GRADIE	NT (FEET/MILE)(10)	1.45	% POOL100) % RIFF	LE0	_%RUN0GF	RADIENT SCORE	8

STREAM: West Fork White River	RIVER MILE	145.1	DATE:	9/21/2004	QHEI SCORE 63
1) SUBSTRATE: (Check ONLY Two Substra TYPE POOL RUN X BLDER/SLAB(10) X BOULDER(9) X COBBLE(8)	POOL X GRAVEL(7) X X SAND(6) X BEDROCK(5) DETRITUS(3) ARTIFIC(0) X <4(0)	RUN SUBS X LIMES X X TILLS SAND SHALI COAL	TRATE ORIGIN (all stone(1) RIP/RAP(0) HARDPAN STONE(0)	SILT CO\ 3) SILT-HEAVY(-2) X SILT-NORM(0)	BSTRATE SCORE 14 /ER (one) SILT-MOD(-1) SILT-FREE(1) edness (check one) MODERATE(-1) NONE(1)
OVERHANGING VEGETATION(1) X ROO	PPOOLS(2) OXBOW TWADS(1) AQUATION	S(1) C MACROPHYTES(1) R WOODY DEBRIS(1)	T (Check only one or 0 EXTENSIVE >7 X MODERATE 25 SPARSE 5-25% NEARLY ABSE	-75%(7) (3)
3) CHANNEL MORPHOLOGY: (Check ONLY SINUOSITY DEVELOPMENT EXCELLENT(7) GOOD(5) X LOW(2) FAIR(3) POOR(1) COMMENTS:	ONE per Category or CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO RECO	X HIGH(MODE	LITY <u>M</u> 3) :RATE(2)	ODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MO	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIAN ZONE AND BANK EROSION: River Right Looking Downstream RIPARIAN WIDTH (per bank) L X (per bank) L X WIDE>150ft.(4) MODERATE 30-150 ft.(3) NARROW 15-30 ft.(2) VERY NARROW 3-15 ft.(1) NONE(0) COMMENTS:	(Check ONE box or Check ONE box or Check ONE box or Check ONE box or Check One of the Check	DDPLAIN QUALIT t per bank) L CROP(0)		BAI L RIAL(0) LD(2) X	RIPARIAN SCORE 6 NK EROSION R (per bank) NONE OR LITTLE(3) X MODERATE(2) HEAVY OR SEVERE(1)
X >4 ft.(6) X POOI 2.4-4 ft.(4) POOI	Y (12) HOLOGY (Check 1) WIDTH-RIFFLE WIDTH(2) WIDTH=RIFFLE WIDTH(1) WIDTH <riffle td="" width(0)<=""><td></td><td>POOL/RUN/RIFFLE TORRENTIAL(-1) FAST(1) MODERATE(1) X SLOW(1)</td><td>CURRENT VELOCITY X EDDIES(1) INTERSTITIAL(</td><td></td></riffle>		POOL/RUN/RIFFLE TORRENTIAL(-1) FAST(1) MODERATE(1) X SLOW(1)	CURRENT VELOCITY X EDDIES(1) INTERSTITIAL(
RIFFLE/RUN DEPTH GENERALLY >4 in. MAX.>20 in.(4) GENERALLY >4 in. MAX.<20 in.(3) GENERALLY 2-4 in.(1) GENERALLY <2 in.(Riffle=0)(0) COMMENTS: 6) GRADIENT (FEET/MILE)(10) 1.14	RIFFLE/RUN SUBSTRA STABLE (e.g., Cobble,B MOD. STABLE (e.g., Pe UNSTABLE (Gravel, Sa NO RIFFLE(0)	ioulder)(2) :a Gravel)(1) nd)(0)	EXTEN MODE LOW(RATE(0) NO	RIFFLE SCORE 0

STREAM: Wes	t Fork Whit	e River	RIVER MILE	156.36	DATE:	9/20/2004	QHEI SCORE 66
TYPE BLDER/SLAB(BOULDER(9) COBBLE(8) HARDPAN(4) MUCK/SILT(2) TOTAL NUMBER OF	POOL F	X X X X X X X X X X	Type Boxes: Chec POOL GRAVEL(7) X SAND(6) X BEDROCK(5) DETRITUS(3) ARTIFIC(0) X <-4(0) ore is based on natural sul	X LIMES X TILLS X SAND SHALL COAL	TRATE ORIGIN STONE(1) RIP/R HARD OSTONE(0)	AP(0) SILT-NORM(0)	UBSTRATE SCORE 14 VER (one) SILT-MOD(-1) SILT-FREE(1) dedness (check one) MODERATE(-1) LOW(1)
2) INSTREAM CO	TY SETATION(1) W WATER)(1)	/PE(Check all X DEEP PORT ROOTW BOULDE	OOLS(2) OXBO	WS(1) TIC MACROPHYTES(1 OR WOODY DEBRIS()	OUNT (Check only one or EXTENSIVE > X MODERATE 2 SPARSE 5-25' NEARLY ABSE	5-75%(7) 6(3)
3) CHANNEL MOR SINUOSITY HIGH(4) MODERATE(3) LOW(2) NONE(1) COMMENTS:	RPHOLOGY: (I DEVELOR EXCELLER GOOD(5) X FAIR(3) POOR(1)	PMENT	DNE per Category of CHANNELIZATION X NONE(6) RECOVERED(4) RECOVERING(3) RECENT OR NO REC	STABI HIGH X MODI LOW	LITY (3) ERATE(2)	MODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MI	IMPOUND ISLAND LEVEED BANK SHAPING
4) RIPARIAN ZON River Right Looking RIPARIAN WIDTH L R (per bank) X WIDE>150ft.(4 X X MODERATE 31 NARROW 15-3 VERY NARRO NONE(0) COMMENTS:	g Downstream (per bank)) 0-150 ft.(3) 0 ft.(2)	ERC L	heck ONE box or C SION/RUNOFF-FLC R (most predomina FOREST, SWAMP(3) OPEN PASTURE/RO RESID.,PARK,NEW F FENCED PASTURE(1)	DODPLAIN QUALI int per bank) w crop(0) rield(1)		JSTRIAL(0) FIELD(2) AGE(1)	RIPARIAN SCORE 7 NK EROSION R (per bank) NONE OR LITTLE(3) X MODERATE(2) HEAVY OR SEVERE(1)
5) POOL/GLIDE A MAX. DEPTH (CI X >4 ft.(6) 2.4-4 ft.(4) 1.2-2.4 ft.(2) <1.2 ft.(1) <0.6 ft.(Pool=0)(0) COMMENTS:		MORPHO POOL W	(12) DLOGY (Check 1) VIDTH>RIFFLE WIDTH(2) VIDTH=RIFFLE WIDTH(1) VIDTH <riffle td="" width(0)<=""><td></td><td>POOL/RUN/RIFF TORRENTIAL(-1) X FAST(1) X MODERATE(1) X SLOW(1)</td><td>LE CURRENT VELOCIT DEDDIES(1) INTERSTITIAL INTERMITTEN</td><td>(-1)</td></riffle>		POOL/RUN/RIFF TORRENTIAL(-1) X FAST(1) X MODERATE(1) X SLOW(1)	LE CURRENT VELOCIT DEDDIES(1) INTERSTITIAL INTERMITTEN	(-1)
GENERALLY >4 in. N GENERALLY >4 in. N GENERALLY >4 in. N GENERALLY 2-4 in. (COMMENTS:	MAX.>20 in.(4) MAX.<20 in.(3) 1) Riffle=0)(0)	1.20	RIFFLE/RUN SUBSTI STABLE (e.g., Cobble MOD. STABLE (e.g., UNSTABLE (Gravel, S NO RIFFLE(0)	e,Boulder)(2) Pea Gravel)(1)	E) MM	ODERATE(0) X No	RIFFLE SCORE 0 S N(2) D RIFFLE(0) RADIENT SCORE 10

STREAM:	West Fork White	e River	IVER MILE	164.53	DATE:	9/20/2004	QHEI SCORE 65
TYPE BLDE BOU COB HARR MUC TOTAL NUM	RATE: (Check ONLY T POOL R ER/SLAB(10) LDER(9) BLE(8) DPAN(4) K/SILT(2) MBER OF SUBSTRATE TYPE re sludge that originates from	X	RAVEL(7) X X AND(6) X EDROCK(5) ETRITUS(3) RTIFIC(0) X < 4(40)	X LIME X TILL X SAN SHA	STRATE ORIGIN ESTONE(1) RIP/RA	AP(0) SILT-NORM(0) SILT-NORM(0)	UBSTRATE SCORE 14 VER (one) X SILT-MOD(-1) SILT-FREE(1) Idedness (check one) X MODERATE(-1) LOW(1)
UNDERCU X OVERHAN	T BANKS(1) GING VEGETATION(1) S (IN SLOW WATER)(1)	X DEEP POOLS ROOTWADS BOULDERS(1)	OXBOW AQUATION	S(1) C MACROPHYTES(R WOODY DEBRIS	(1)	UNT (Check only one or EXTENSIVE > X MODERATE 2 SPARSE 5-25' NEARLY ABSI	5-75%(7) %(3)
3) CHANNE SINUOSIT HIGH(4) MODERAT X LOW(2) NONE(1)	EXCELLEN GOOD(5) FAIR(3) X POOR(1)	MENT CHA	per Category or INNELIZATION ONE(6) ECOVERED(4) ECOVERING(3) ECENT OR NO RECC	STAE HIGH X MOD	H(3) DERATE(2)	MODIFICATION/OTE SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL M	IMPOUND ISLAND LEVEED BANK SHAPING
River Right RIPARIAN L R (pei X X WIDE MOD		EROSIO L R (N/RUNOFF-FLOC MOST Predominant OREST, SWAMP(3) PEN PASTURE/ROW ESID.,PARK,NEW FIE ENCED PASTURE(1)	DDPLAIN QUAL t per bank) L CROP(0)		B <u>A</u> L STRIAL(0) FIELD(2) X GE(1)	RIPARIAN SCORE 8 ANK EROSION R (per bank) NONE OR LITTLE(3) X MODERATE(2) HEAVY OR SEVERE(1)
5) POOL/G	LIDE AND RIFFLE/RU PTH (Check 1)	MORPHOLOG POOL WIDTH		O POOL = 0	POOL/RUN/RIFFI TORRENTIAL(-1) X FAST(1) X MODERATE(1) X SLOW(1)	LE CURRENT VELOCIT EDDIES(1) INTERSTITIAL INTERMITTEN	.(-1)
GENERALI GENERALI GENERALI COMMEN	Y >4 in. MAX.>20 in.(4) Y >4 in. MAX.<20 in.(3) Y 2-4 in.(1) Y <2 in.(Riffle=0)(0)	S M U N	TABLE (e.g., Cobble,E IOD. STABLE (e.g., Pe NSTABLE (Gravel, Sa O RIFFLE(0)	Boulder)(2) ea Gravel)(1)	EX MC SP	DDERATE(0) X N	RIFFLE SCORE 0 SS W(2) O RIFFLE(0)

STREAM:	West Fork White I	River RIVER MILE	194.09	DATE:	9/28/2004	QHEI SCORE	66
TYPE BLDE BOUL COBI HARRI MUC TOTAL NUM	POOL RIFF ER/SLAB(10) LDER(9) BLE(8) DPAN(4) K/SILT(2) MBER OF SUBSTRATE TYPES: re sludge that originates from po	X GRAVEL(7)	OL RIFFLE SUB: X LIME X TILL: X SAN SHA COA	STRATE ORIGIN (a	SILT CO SILT-HEAVY(-2) N(0) X SILT-NORM(0)	BSTRATE SCORE VER (one) SILT-MOD(-1) SILT-FREE(1) edness (check one) X MODERATE(-1) LOW(1)	14
UNDERCU X OVERHAN	T BANKS(1) GING VEGETATION(1) S (IN SLOW WATER)(1)	ROOTWADS(1)	BOWS(1) UATIC MACROPHYTES(3S OR WOODY DEBRIS	1)	NT (Check only one or (EXTENSIVE >7 X MODERATE 25 SPARSE 5-25 NEARLY ABSE	5%(11) -75%(7) 6(3)	_
SINUOSITY HIGH(4) MODERAT X LOW(2) NONE(1)	E(3) DEVELOPME EXCELLENT(7) GOOD(5) X FAIR(3) POOR(1)		STAB HIGH X MOD LOW	BILITY H(3) DERATE(2) //(1)	MODIFICATION/OTH SNAGGING RELOCATION CANOPY REMOVAL DREDGING ONE SIDE CHANNEL MO -WEll field.	IMPOUND ISLAND X LEVEED BANK SHAPING	13
River Right RIPARIAN L R (per WIDE X X MOD NARI	Looking Downstream WIDTH (per bank) bank) =>150ft.(4) ERATE 30-150 ft.(3) ROW 15-30 ft.(2) Y NARROW 3-15 ft.(1) E(0)	EROSION/RUNOFF-F L R (most predomin FOREST, SWAMP OPEN PASTURE/F X RESID.,PARK,NEV FENCED PASTUR	LOODPLAIN QUAL nant per bank) L (3) ROW CROP(0) V FIELD(1)		TRIAL(0) ELD(2) X E(1)	RIPARIAN SCORE NK EROSION R (per bank) NONE OR LITTLE(: X MODERATE(2) HEAVY OR SEVER	3)
,) bl=0)(0)	QUALITY (12) MORPHOLOGY (Check 1) POOL WIDTH-RIFFLE WIDTH POOL WIDTH-RIFFLE WIDTH POOL WIDTH-RIFFLE WIDTH	(1)	POOL/RUN/RIFFL TORRENTIAL(-1) X FAST(1) MODERATE(1) SLOW(1)	E CURRENT VELOCITY EDDIES(1) INTERSTITIAL(-	-1)	
GENERALI GENERALI GENERALI COMMEN	_Y >4 in. MAX.>20 in.(4) _Y >4 in. MAX.<20 in.(3) _Y 2-4 in.(1) _Y <2 in.(Riffle=0)(0)	RIFFLE/RUN SUBS STABLE (e.g., Cob MOD. STABLE (e.g. UNSTABLE (Grave NO RIFFLE(0)	ble,Boulder)(2) g., Pea Gravel)(1)	EXTI MOD SPA	DERATE(0) X NC	RIFFLE SCORE V(2) ORIFFLE(0) ADJENT SCORE	